

Heterogeneity of outcomes in randomized controlled trials on implant prosthodontic therapy is hindering comparative effectiveness research: meta-research study

Vardić, Ante; Puljak, Livia; Galić, Tea; Viskiće, Joško; Kuliš, Ena; Poklepović Peričić, Tina

Source / Izvornik: **BMC Oral Health, 2023, 23**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1186/s12903-023-03658-9>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:127:413399>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-11**



Repository / Repozitorij:

[University of Zagreb School of Dental Medicine
Repository](#)



RESEARCH

Open Access



Heterogeneity of outcomes in randomized controlled trials on implant prosthodontic therapy is hindering comparative effectiveness research: meta-research study

Ante Vardić¹, Livia Puljak^{2*} , Tea Galić³, Joško Viskić⁴, Ena Kuliš¹ and Tina Poklepović Peričić³

Abstract

Background Consistency in outcomes across clinical trials allows for comparing and combining results from different studies. A core outcome set (COS), representing a minimally agreed standardized group of outcomes that should be monitored and measured through research in a specific field of medicine, is not yet available for trials in implant prosthodontic (dental implant) therapy. This meta-research study aimed to analyze outcomes used in clinical trials on implant prosthodontic therapy.

Methods We searched the Cochrane Oral Health Group (COHG) register to identify systematic reviews of interventions in implant prosthodontic therapy published by October 2023. From the randomized controlled trials (RCTs) included in the relevant reviews, we extracted data on the characteristics of the included trials and the outcomes used. We categorized outcomes into domains.

Results From 182 systematic reviews in the COHG register, we included 11 systematic reviews on dental implant therapy. The reviews included 117 unique RCTs with 4725 participants, published from 1995 to 2020, which analyzed 74 different outcomes. Using different definitions, implant failure was analyzed in 73 RCTs. Seventeen RCTs did not define implant failure. Failure was most often (30 RCTs) followed up for one year. Only one RCT assessed implant failure after five years. Trials used 17 definitions of implant failure, while 17 trials did not report on the criteria of implant failure. Complications were analyzed in 48 RCTs, although they were not clearly defined in 12 RCTs. Failure of prosthodontic supra-structure was analyzed in 74 RCTs, with definitions of failure and criteria not clearly defined in 44 RCTs. Trials considered adverse events, peri-implant tissue health, patient attitudes, and other outcomes, including cost, aesthetics, or procedure duration. These outcomes were often different between trials. Twenty-six outcomes were used only once per study.

Conclusions Clinical trials in implant prosthodontics used different outcomes, different definitions of outcomes and used different times to monitor them. Standardization of outcomes is necessary to allow comparability and evidence synthesis about the effectiveness of implant prosthodontic therapy.

*Correspondence:

Livia Puljak

livia.puljak@unicath.hr; livia.puljak@gmail.com

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords Core outcomes, Implant prosthodontics, Randomized controlled trials, Meta-research

Background

Implant prosthodontics includes treatment planning and restoration of dental implants to replace a lost tooth. Dental implants can restore masticatory function and improve quality of life [1–3], and have therefore become a popular treatment option for partially or completely edentulous patients [4]. However, despite the high success rate and low percentage of complications of such therapy, some patients experience dental implant failure [5].

Depending on when it occurs, implant failure may be classified as early or late, i.e. before or after functional implant loading [5]. Early failure represents the inability to establish osseointegration and includes biological complications affecting the hard and soft tissues surrounding the implant [6–8]. Occlusal forces during chewing, swallowing and biting or during lateral movements of the jaw significantly affect the dental implant-retained prosthesis [9]. Late failure is defined as the inability to maintain the established osseointegration and implant function. It is accompanied by biological and mechanical complications, such as fracture of the implant body, screw body, or implant supra-structure [5, 6, 10].

However, there is no single clear definition of the criteria for assessing the failure of implant therapy [11].

Criteria to assess the success of dental implant therapy have changed over time. The most frequently considered criteria have been survival of dental implants, stability of prosthesis, radiographic evidence of bone loss, and the absence of infection of the peri-implant tissues [12–14]. Based on the available criteria, dental implant therapy is considered successful if there is no mobility of the implant at the start of the prosthetic phase, bone loss is less than 0.2 mm per year after the first year, there is no radiolucency around the implant, there are no signs of peri-implantitis with suppuration, and no symptoms of pain, neuropathy or nerve paraesthesia [12, 15, 16].

The appearance of the soft tissues surrounding the implant, assessment of prosthodontic supra-structure, aesthetics and patient satisfaction were also proposed as criteria of success [17–19].

Heterogeneity in defining the success and failure of dental treatment therapy in clinical practice indicates the possibility that clinical studies may use different outcomes [12–14]. Furthermore, inconsistent outcomes across studies make it impossible to directly compare and systematically summarize all available evidence by combining results from various studies [12, 15, 17]. Thus, it is crucial to standardize outcomes and outcome measures in research on dental implant therapy.

Ideally, a core outcome set (COS) should be used, representing a minimally agreed standardized group of outcomes that should be monitored and measured through research in a specific field of medicine to enable comparison and combination of the results from different studies [20]. However, in the field of dental implant therapy, there is currently no COS, and so far, no studies have examined all outcomes used in clinical trials on implant prosthodontic treatment.

Therefore, this study aimed to map all the outcomes assessed in clinical studies about the efficacy of dental implant therapy and analyze outcomes used to define treatment success or failure of dental implants.

Materials and methods

Study protocol

We developed a protocol for this study before the study started. The protocol was not published; it is available in Supplementary file 1.

Study design

We conducted a cross-sectional meta-research study of randomized controlled trials (RCTs) included in published systematic reviews (SRs) from the Cochrane Oral Health Group (COHG) register. The COHG utilizes Cochrane Oral Health's Trials Register, which conducts regular searches in the databases CENTRAL, MEDLINE and Embase.

Eligibility criteria

We analyzed RCTs included in SRs of interventions assessing the effectiveness of different treatment approaches to dental implant therapy, including different implant techniques, different sizes and shapes of implants, or different times for implant loading. We also included SRs that analyzed preoperative therapy before implantation.

Search

We used the COHG register of published SRs (available at: <https://oralhealth.cochrane.org/oral-health-evidence>) to identify eligible SRs. The following search terms were used: “dental implants”[All Fields] OR “dental implants”[MeSH Terms] OR „dental implant”[Text Word] OR „oral implantation”[Text Word] OR „oral implanting”[Text Word] OR „dental implanting”[Text Word].

We screened 182 published SRs from the COHG register, which covered a wide range of dental medicine topics. The search was first conducted on May 27th 2022 and then updated on October 21st 2023.

Screening

Two authors (AV, EK) independently screened titles and abstracts of all SRs from the COHG list of reviews in the first screening phase. Subsequently, in the second screening phase, two authors (AV, EK) independently assessed full texts of potentially eligible SRs for inclusion. Disagreements about including full texts were resolved by discussion between the two authors (AV, EK) or by consulting the senior author (TPP).

Data extraction

Initially, full texts of RCTs included in the eligible SRs were obtained. A data extraction form was developed for this study and was first piloted on three randomly selected SRs. After final refinements to the extraction form, one author (AV) extracted the data, and another (EK) verified the extractions. The following data were extracted: title of the SR, first author of the SR, publication year of the SR, the list of all included RCTs, first author of the RCT, publication year, number of participants, participants' health status, the country in which the RCT was conducted, and the list of all outcomes as they were reported in the [Results](#) section of each RCT.

Additionally, from each RCT, we extracted information concerning outcomes related to the implant or prosthetic supra-structure, including definitions of the implant or prosthetic failure and follow-up times, and the type of prosthodontic supra-structure used (fixed or removable). Also, data were extracted on outcomes related to the status of the tissues surrounding the implant, as well as specific outcomes considered as postoperative complications and adverse events. All other outcomes, such as those related to how patients accept dental implants, clinicians' preferences towards a specific procedure, evaluation of aesthetics, and difficulty or duration of the procedure, were also extracted and included in analyses.

Outcome categorization

Outcomes were categorized according to their similar features into the following domains: outcomes related to the implant itself or the prosthetic supra-structure, outcomes related to complications and adverse events, outcomes related to peri-implant tissue health, and patient-related outcomes. The domain concerning the health of the peri-implant tissues was further subcategorized into oral hygiene outcomes, outcomes related to soft tissue, and bone-related outcomes. Finally, the remaining outcomes that did not fit in any of the previous four domains were categorized as "other outcomes", e.g. cost of therapy, treatment time, etc.

The decision on categorizing domains was based on the discussion among authors (AV, TG, JV, TPP). Data were entered into a Microsoft Office Excel spreadsheet and appropriately coded.

Data analysis

We used descriptive summary statistics with absolute numbers and percentages to present the number of studies in each SR, the number of study participants, and the frequency of each outcome across the studies. Data were analyzed using MedCalc, version 19.4 (MedCalc Software, Ostend, Belgium).

Results

General characteristics of included studies

Among the 182 screened SRs, 168 records that did not fit the inclusion criteria during the first screening phase were excluded. The remaining 14 SRs were reviews that were in any kind of way associated with the implant prosthodontic therapy. However, after analyzing the full text of those 14 SRs, we excluded three more SRs, of which two were excluded because dental implant therapy for replacing missing teeth was not the intervention of interest. One of the mentioned reviews instead assessed mini-implants' use for orthodontic anchorage [21], while the other review assessed the effects of various interventions, hyperbaric oxygen therapy and antibiotics, for preventing osteoradionecrosis in the jaws of patients treated for head and neck cancer before implant placement [22]. The third review was excluded [23] because it did not include a single RCT, i.e. it was an empty review.

Finally, 11 SRs [24–34] were included, which were published between 2009 and 2021. The selected SRs included a total of 118 RCTs. Among the 118 RCTs, there was one overlapping trial [35]. Thus, we analyzed 117 unique RCTs, which were published from 1995 to 2020.

We used the PRISMA flowchart to depict the flow of SRs during the searching and screening phase of this study [36]. A list of all included SRs and RCTs is provided in Supplementary file 2. The flow diagram of screening and study inclusion is shown in Fig. 1.

The 117 RCTs were conducted in 27 countries worldwide; most trials (N=40) took place in Italy. For five trials, it was not reported where the trials were conducted, and one was a multicenter trial reportedly conducted in ten countries, but with no information on which countries. The list of countries with the number of trials conducted in each country is available in Supplementary file 3.

The 117 RCTs included a total of 4725 participants (range: from 5 to 496 participants). In 92 RCTs, the participants were healthy individuals; those studies excluded participants whose health conditions could affect the success of the therapy or cause a deviation in the results. In two RCTs, participants had an unspecified periodontal disease but were otherwise healthy. In one RCT, all participants had head or neck cancer in their medical history and were treated with radiotherapy. One study included participants using long-term non-steroidal




Fig. 1 Flow diagram. Acronyms: COHG=Cochrane Oral Health Group; SR=systematic review

Table 1 Outcomes related to the implant itself or prosthetic supra-structure

OUTCOME	N of trials
Implant failure	73
Supra-structure failure	74
Implant Stability Quotient (Osstell device)	11
Implant survival	10
Implant success	3
Implant mobility (periostest)	8
Crown success	2

anti-inflammatory drugs (NSAID) therapy, requiring antibiotic prophylaxis before surgical procedures. 22 RCTs did not specify the health condition of the subjects.

Supplementary file 4 consists of raw data extracted from the 117 RCTs.

Analysis of outcomes

A total of 74 different outcomes were used in 117 RCTs. Six RCTs [37–42] assessed only one outcome, and four RCTs [43–46] assessed just two outcomes. Outcomes were categorized into five main domains, as well as ‘other outcomes’.

Outcomes related to the implant itself or prosthetic supra-structure

We identified seven outcomes related to the implant itself or prosthetic supra-structure (Table 1). Implant failure was assessed in 73 different trials [16, 35, 46–116] (Table 1).

Ten trials [63, 64, 117–124] assessed implant survival, of which only four trials reported on the criteria for defining survival. Si et al. [63] referred to the definitions suggested by Buser in 1997 [14] and Cochran in 2002 [125] that consider implant mobility, pain and discomfort, peri-implant infection, and continuous radiolucency around the implant. According to these definitions, implant loss, mobility or removals in case of progressive marginal bone loss, severe peri-implant infection, or implant fracture were considered implant failure. Torres et al. [64] defined survival as implants remaining in situ during clinical observations, while Patel et al. [124] reported survivals as “any dental implant which remained in situ without signs of mobility”. In the trial by Enkling et al. [119] implant was considered survival if it remained stable when tested by removal resistance. No other information was provided on how this testing was done.

Six trials [118, 119, 121–124] reported data on survival after a one-year follow-up, while three studies al [63, 64, 117] provided a two-year follow-up for survival. Horwitz et al. [120] assessed survival after three months.

Table 2 Criteria of implant failure from 73 trials that used this outcome

DEFINITION CRITERIA (verbatim when possible)	N of trials
Criteria by Albrektsson et al. 1986	10
"..implant mobility and removal of stable implants dictated by progressive marginal bone loss or infection."	8
Loss of implant	6
"..presence of any mobility of the individual implant and/or any infection."	6
Implant mobility	5
Lack of osseointegration	5
Mobility and/or any situation dictating removal	3
"..implant mobility and removal of stable implants dictated by progressive marginal bone loss, infection or implant fracture."	3
"..lack of implant stability, presence of radiolucent zone around the implants, mucosal suppuration and pain."	2
"..implant mobility, removal of stable implants dictated by progressive marginal bone loss or infection, and any mechanical complications (e.g. implant fracture) rendering the implant unusable. Also, implants that had to be removed at implant insertion due to lack of stability and the risk of falling into the sinus cavity."	1
"implant mobility, removal of stable implants as a result of progressive bone loss and implant fracture"	1
"A failed implant was defined as the presence of signs of infection and/or radiographic peri-implant radiolucencies that could not respond to a course of antibiotics and/or judged a failure after performing an explorative flap surgery by an experienced periodontologist."	1
Mobility, bone loss, radiolucency, pain, discomfort and/or neurosensory alteration	1
"..fracture, mobility when tested, peri-implant radiolucency, pain, discomfort, infection, and/or marginal bone loss that could not be alleviated by clinical intervention."	1
"..radiolucency around the implant, mobility and suppuration, pain or ongoing pathologic processes."	1
"..implant mobility; presence of peri-implant radiolucency, recurrent peri-implant infection, continuous or recurrent pain, or structural failure of the implant; and > 0.2 mm bone resorption between any two visits."	1
"..progressive peri-implant loss of bone that exceeds the limits of tolerable bone absorption after successful osseointegration."	1
Definition not reported	17

Various definitions of implant failure were reported, with as many as 17 different criteria to define failure across trials (Table 2). Most trials used the criteria proposed by Albrektsson et al. in 1986 [11] that consider implant mobility, pain, dysesthesia, peri-implant infection with suppuration, peri-implant radiolucency, and peri-implant bone resorption of more than 1.5 mm in the first year of function and more than 0.2 mm in the following years (Table 2). In 17 trials [35, 67, 68, 80, 86, 95, 97, 101, 104, 105, 108, 110–115], the definition of failure was not provided.

Furthermore, trials used 13 different follow-up times to assess the failure of implant therapy (Table 3). The most commonly used follow-up time was 12 months, followed

Table 3 Follow-up time for implant failure in 73 trials that used this outcome

FOLLOW-UP TIME	N of trials
3 months	2
3–4 months	2
4 months	3
5 months	2
6 months	10
9 months	1
1 year (12 months)	30
18 months	2
19 months	1
2 years (24 months)	11
3 years (36 months)	7
4 years (48 months)	1
5 years (60 months)*	1

*recommended follow-up time for implant failure (by Albrektsson et al., 1986)

by 24 months. Only one trial [52] provided data after a five-year follow-up, which was recommended for assessing implant failure by Albrektsson et al. in 1986 [11]. Almost a quarter of trials (23%) followed-up implant failure for more than 12 months (Table 3).

Among 74 trials that assessed prosthesis failure [35, 43–46, 49, 50, 52–99, 101–106, 117–119, 121–123, 126–131], two types of prosthetic supra-structure were used: fixed (screwed or cemented) (N=46; 62%) and removable (overdenture) (N=14; 18%). In eight (11%) trials [49, 50, 63, 64, 101, 104–106], the type of prosthetic supra-structure was not reported. Different definitions of prosthesis failure outcome were used in 30 trials [43–45, 49, 50, 52–60, 62, 72–74, 77, 79, 81, 82, 87, 92, 93, 98, 102, 128–130], but mostly considered prosthesis failure as the inability to place planned prosthesis due to implant failure or loss of prosthesis following implant failure [49, 50, 52, 53, 55–59, 62, 72–74, 92, 93].

Hall et al. [81] assessed the failure of fixed prosthetic supra-structures using criteria proposed by Walton [55], and two other trials published by Felice et al. in 2009 [56, 57] evaluated the success of mandibular overdentures using the six-step protocol proposed by Payne [58]. The criteria to assess prosthodontic success for implant retained overdentures referred to in these trials [56, 57, 81] consider the following: patrix and matrix loosening, fracture, number of times they were activated or replaced, fracture of implant overdenture, the need to reline or construct a new overdenture, and peri-implant or inter-abutment mucosae enlargement. These criteria were first included in a classification protocol proposed by Walton in 1998 [55]. In 2002 Payne [58] published a protocol for prosthodontic maintenance, including the same specific categories. Therefore, we considered these criteria as one outcome in our analyses. The trials [56, 57, 81] that assessed prosthodontic maintenance using

general categorization according to either the classification protocol proposed by Walton [55], or the Payne protocol [58] were therefore presented together.

A total of 44 trials [35, 61, 63–71, 75–78, 80, 16, 83–86, 88–91, 94–97, 99, 101, 103–106, 117–119, 121–123, 126–128] out of 74 trials did not report any details about defining this outcome (Table 4).

Outcomes related to postoperative complications and adverse events

From 117 RCTs, only 14 trials [47–51, 107, 108, 110–115, 132] referred to outcomes related to adverse events. Postoperative pain [51] and postoperative infection [132] were assessed as adverse events in one trial each. A set of outcomes, including erythema multiforme, urticaria, nausea, vomiting, diarrhoea etc., were assessed in two trials [49, 50]. However, most trials (N=10) reported on “adverse events” using a general term without specifying any details about the outcome.

Postoperative complications were assessed in 48 trials [35, 48–51, 53–62, 64, 65, 69, 72–75, 82, 87, 92–94, 96, 98, 99, 101, 102, 104–108, 110–114, 119, 122, 123, 127, 128, 133], but they used different sets of outcomes to assess complications (Table 5). Most trials (N=16) [53, 55–60, 62, 72–75, 92, 93, 106, 133] considered any prosthetic or biological complications like wound or implant infection, mucositis, abscesses or periimplantitis. One trial [96] reported complications as “any biologic complications”, while some trials assessed more specific outcomes, like edema, erythema, wound dehiscence, inflammation etc.

Two trials [102, 108] specified complications as peri-implant mucositis and peri-implantitis. Criteria proposed by Enislidis [134] were used in one trial [94], while other trials (N=16) [48–51, 54, 64, 65, 82, 98, 99, 101, 105, 113, 114, 122, 123] included outcomes like dehiscence, occlusal interference, tilting of segment, pain, swelling, hypesthesia, fracture of the basal bone, breakage of distractor, infection, inflammation, or breakage either in terms of mechanical block preventing distractor activation, instability of distractor or disengagement of a threaded rod from the basal stabilizing plate, as well as fracture of transport segment.

The remaining 12 trials [35, 61, 69, 87, 104, 107, 110–112, 119, 127, 128] did not specify any details about complications.

Eleven trials [48–51, 107, 108, 110–114] considered both types of outcomes, postoperative complications and adverse events. However, there were some overlaps between these two groups of outcomes. Namely, postoperative pain and infection were considered postoperative complications in some trials [54, 99, 101, 105], while in others, they were regarded as adverse events [51, 132].

Table 4 Definitions of prosthesis failure from 74 trials that used this outcome

DEFINITION	N of trials
Prosthesis that could not be placed due to implant failure, or loss of prosthesis secondary to implant failure	15
Prosthetic maintenance by general categorization, including criteria by Walton 1998 and Payne 2002s	7
Prosthesis could not be placed due to implant failure, or loss of prosthesis secondary to implant failure, or any prosthesis in need of replacement	2
Fracture or component failure	3
Prosthesis mobility	2
Need for adjustments of repair	1
Definition not reported	44

Table 5 Outcomes reported as postoperative complications

OUTCOME (Verbatim where appropriate)	N of trials
Any prosthetic or biological complications	16
Postoperative pain or any kind of discomfort	5
Peri-implant mucositis or peri-implantitis	2
“..wound dehiscence, suppuration, fistula, abscess, osteomyelitis, etc.”	2
Any biologic complications	1
Not specified	12
“..internal and external edema, internal and external erythema, pain, heat, and exudate.”	1
“Inflammation, redness of the mucosa, wound dehiscence, sequestration, and loss of bone particles.”	1
Sinus membrane dehiscence	1
“...such as unexpected deviations from the normal treatment outcome; examples of biological complications are haemorrhaging during and after implant placement and/or peri-implantitis.”	1
“Minor” or “major” complications based on the criteria established by Enislidis et al.	1
“..inflammation, wound infection, wound dehiscence, sensory disturbances of lip and chin.”	1
“..pain, fatigue; bowel function, breathing, appetite and sleep disorders.”	1
Swelling	1
Allergic reactions, swellings, abscesses or infections	1
“Post-operative swelling, bruising, suppuration and wound dehiscence”	1

Indicators of peri-implant tissues health

Trials used various outcomes to assess the status of the tissues surrounding implants which we categorized into three subdomains: outcomes related to oral hygiene, outcomes related to soft tissue, and bone-related outcomes (Table 6). The plaque index was most often (N=22) [54, 61, 70, 79, 80, 83, 99, 100, 105, 110, 115, 116, 119, 123, 135–141] used to assess the level of oral hygiene around implants. There were 16 different outcomes related to the soft tissue surrounding implants, of which the most common outcomes were “probing pocket depth” (N=37)

Table 6 Indicators of peri-implant tissue health

SUBDOMAIN	OUTCOME	N of trials	
OUTCOMES RELATED TO ORAL HYGIENE	Plaque index	22	
	Plaque accumulation	16	
	Calculus (yes/no)	3	
	Lobene stain index	2	
	OUTCOMES RELATED to SOFT TISSUE	Probing pocket depth	37
		Bleeding on probing index	28
		Gingival recession	10
		Gingival index	9
		Sulcus bleeding index	8
		Clinical "attachment" level	8
BONE-RELATED OUTCOMES	Width of the keratinized mucosa	8	
	Microbiological evaluation	7	
	Mucosa level	5	
	Soft tissue thickness	3	
	Papilla index	3	
	Papilla levels	2	
	Bleeding time index	1	
	Crevicular fluid flow rate	1	
	Soft tissue volumetric analysis	1	
	Thickness of buccal wall	1	
	Radiographic peri-implant marginal bone level changes	32	
	Height and width of alveolar ridge	15	
	Peri-implant marginal bone levels	14	
	Vertical bone gain after augmentation	11	
	Peri-implant marginal bone level changes	7	
	Radiographic bone gain	4	
	Bone gain maintenance over time	1	
Bone level changes after loading	1		
Bone to implant distance	1		
Vertical distraction distance	1		
Bone width at implant site	1		
Maxillary bone width changes over time	1		

[46, 54, 61, 69–71, 76, 78, 80, 83, 84, 99, 100, 103, 105, 107–115, 118, 119, 123, 124, 128, 133, 135, 138, 139, 141–143], and "bleeding on probing index" (N=28) [46, 54, 61, 76, 99, 103, 105, 107–116, 118, 119, 123, 133, 135, 136, 138, 134, 141–143]. Trials used 12 different bone-related outcomes. The most commonly used outcomes were "Radiographic peri-implant marginal bone level changes" (N=32) [46, 60–62, 66, 69–71, 73, 74, 84–88, 90, 91, 99, 106, 118, 121, 123, 127, 128, 143], "height and width of the alveolar ridge" (N=15) [37–42, 84, 105, 106, 116, 124, 129, 131, 144, 145] and "peri-implant marginal bone levels" (N=14) [52, 55–59, 63, 75, 92, 93, 108, 119, 128, 133]. We extracted data regarding the type of radiographic imaging used to evaluate peri-implant marginal bone level changes and found that most trials (N=20) [46, 69–71, 73, 74, 84–88, 106, 118, 127, 128] reported using standardized intraoral imaging. Periapical imaging was used in eight trials [60–62, 66, 77, 90, 91, 121], while

Table 7 Patient-related outcomes

OUTCOME	N of trials
Patient satisfaction	19
Patient preference	9
Prosthetics acceptance	2
Patient compliance	1
Anxiety during treatment	1
Functional assessment and quality of life	1
Subjective chewing ability	1
Subjective evaluation of taste and change in taste	1

Table 8 Other outcomes

OUTCOME	N of trials
Histomorphometric evaluation	13
Aesthetics	10
Preoperative augmentation failure	8
Duration of the operative procedure	7
Need for additional augmentation	4
Treatment time (defined as the time from the start of preoperative procedures to the placement of prosthetic supra-structure)	3
Cost of therapy	2
Clinician preference	2
Implant percussion	1
Days needed to start the prosthetic rehabilitation	1
Difficulty of the procedure (in a technical sense)	1

panoramic [76, 143] and extra-oral oblique lateral imaging [99, 123] were used in two trials each.

Patient-related outcomes

Trials used eight different patient-related outcomes (Table 7). Patient satisfaction (N=19) [43–45, 56, 58, 69, 72–74, 99, 101, 106, 119–121, 123, 128, 130, 131] and patient preference (N=9) [53–57, 92, 102, 106, 130] were used most commonly across trials.

Other outcomes

The remaining outcomes could not be categorized into any of the previous groups and were categorized as "other outcomes" (Table 8). Histomorphometric evaluation (N=13) [35, 61, 64, 65, 92, 95, 97, 98, 105, 106, 126, 131, 144] was the most commonly used outcome across trials. It represents the histological evaluation of specimens of bone tissue to assess either newly formed bone tissue, bone graft material, loose connective tissue, quality of the bone, or the presence of bone resorption, etc. Only one trial provided no information about the process of histomorphometric evaluation [98], while the remaining 12 trials reported how the process was carried out.

Aesthetics was assessed in ten trials [56, 68, 69, 105, 106, 117, 118, 121, 127, 128], of which in three RCTs [56, 117, 127], aesthetics was evaluated by the dentists, and in one trial [118] it was evaluated by patients. In three trials [68, 121, 128], aesthetics was evaluated by both dentists

and patients. One RCT [69] used an independent blinded evaluator. Two RCTs [105, 106] used objective evaluation using validated indices to assess aesthetics. One trial [105] used colour blending of the grafted site with the adjacent soft tissues and the other trial [106] used pink aesthetic scores (PES). Two trials assessed clinicians' preferences that refer to either operator's preferences regarding different techniques applied at implant placement [53], or preferences towards augmentation procedures before implant placement [57].

Discussion

This meta-research study found considerable heterogeneity in outcomes used in RCTs on implant prosthodontics. Overall, 74 different outcomes were used in 117 RCTs, of which 24 were used in one trial only. Trials used a wide range of different criteria to define the outcomes. Most trials provided no information on the criteria used to define these specific outcomes. Furthermore, trials measured outcomes at very different times.

With the wide use of implants to replace lost teeth [1], there is a parallel increase in the number and the extent of research in the field of dental implant therapy [146]. However, our study indicates the need for improvement in the consistency and usability of outcomes used in the field. The most commonly assessed outcomes included implant and prosthesis failure, postoperative complications, adverse events, and implant survival. For implant failure, the most commonly used outcome, 17 different definitions were used. Researchers used 13 different follow-up times for implant failure, most commonly one year period, reported in 30 studies. Only one trial analyzed implant failure at five years, which is the recommended minimum time for monitoring the success rate of implant prosthetic therapy according to the criteria of Albrektsson et al. [11].

The short-term follow-up can be understood as a pragmatic choice of trialists. RCTs are complex studies requiring intense resources – time-consuming, expensive, and often burdensome on patients [147]. However, an inadequate trial follow-up period contributes to a lack of reliable evidence for decision-makers [148]. Long-term data collection and analysis are crucial when evaluating a procedure such as dental implant placement [149].

Furthermore, adequate outcomes need to be chosen to evaluate the success/ failure of dental implants properly. Considering the rate of complications in implant-supported fixed partial dentures (FPD) after five years, it is important to include prosthesis success in analyses of the overall success of dental implants [150]. Namely, studies should evaluate a long-term primary outcome by considering the implant prosthetic complex as a whole [151].

In our study, survival, prosthetic failure, and implant failure were assessed separately in many studies. A

systematic review by Papaspyridakos et al., which was published in 2012 [151], addressed success criteria in RCTs and prospective studies on implant dentistry published from 1980 to 2010. They pointed out that in the dental implant literature, survival/success rates of single parameters were often presented, but that single parameters used as success criteria should be regarded as surrogate endpoints, as they are often used to compensate for the lack of well-defined primary outcomes. They advised that, for example, bone loss or any other outcome alone within an implant prosthodontic rehabilitation should not be considered as the measure of success. Papaspyridakos et al. argued that current advances in contemporary implant prosthodontics, coupled with high patients' expectations, necessitate a more comprehensive definition of success criteria for implant/prosthodontic procedures. They suggested that future studies should choose outcomes that reflect the complexity of the implant-prosthetic complex and consider multiple outcomes as measures of success or implant failure [151].

Papaspyridakos et al. found similar results to ours, with different trials using different criteria to assess the success of the implant therapy. Furthermore, they found that the reported success rate of dental implant therapy consistently decreased when the number of parameters included for measuring success was increased [151].

In 2009, Gallucci et al. proposed success criteria for implant-supported fixed complete dental prostheses (FCDPs) based on the implant, peri-implant tissues, prosthodontic, and subjective parameters. They suggested that complete dental prostheses (CDPs) were deemed as successful when a total of four or fewer complications (mild or moderate severity) occurred and when these could be addressed chair-side in a single visit [152].

Our study showed that patient satisfaction was analyzed in only 19 of 117 trials (16%). According to Levi et al., patient satisfaction with overall treatment should be rated good or excellent for the treatment outcome to be considered successful [153].

Furthermore, a conceptual framework proposed for understanding the outcomes of dental implant therapy also includes psychological outcomes related to the patient and economic aspects. However, in our sample, only two trials assessed the cost of treatment [154].

Aesthetics is also an important outcome in dental implant treatment for both patients and clinicians [155]. The Pink Esthetic Score (PES) and the White Esthetic Score (WES) have been proposed for measuring aesthetics in implant prosthodontics [16, 18]. In this study, only ten trials assessed aesthetics. However, only two trials used validated scales to assess aesthetics, the colour blending of the grafted site with the adjacent soft tissues and the PES, while eight trials did not explain in detail how they assessed aesthetics at all. Those studies only

mentioned that aesthetics was assessed by the clinician or patient.

Problems with outcome heterogeneity, different definitions, and lack of definitions were noted in other research fields as well. For example, a systematic review found ten different definitions for postoperative mortality in esophageal cancer research, most of which were not clearly described, and there were different interpretations of the term in-hospital mortality [156, 157].

Heterogeneity in outcomes measured across studies in the same disease or treatment hamper clinical evaluation, trial comparability, and effective evidence synthesis [157]. The results of heterogeneous studies cannot be combined, compared in systematic reviews, and further used for developing clinical practice guidelines. Therefore, their applicability in the context of other research and clinical practice remains questionable, including their contribution to overall scientific knowledge and clinical practice as well.

For example, 24 out of 74 outcomes found in analyzed trials were used once per trial. A meta-analysis of such outcomes is not possible. In addition, another 14 outcomes were measured in five or fewer primary studies, meaning that overall more than half (38 of 74) of the outcomes were replicated in more than 5% of studies. If we were to look only at outcomes that were repeated in more than 10% of the primary studies, we would have only 13 such outcomes. A deleterious example of single use of outcomes was reported in a newly published study from the field of oncology that found as many as 25,000 different outcomes used only once in oncology trials [158].

Furthermore, for clinically meaningful outcomes, all clinical trials of interventions must analyze not only efficacy but also harms. In our study, only 14 trials out of 114 analyzed adverse events.

Adverse events are less likely to be reported than efficacy outcomes, and different methods of assessing adverse effects produce different reported incidences [159, 160], despite available reporting guidelines suggesting the inclusion of adverse events in research papers [161, 162]. Golder et al. found that a median of 43% of published studies reported adverse events data, compared with a median of 83% of unpublished studies. A wider range of specific adverse events was found in sources other than published journal articles. In addition, when published and unpublished reports of the same study were compared, it was shown that the unpublished version was more likely to contain adverse effects data (median 95%) compared with the published version (median 46%) [163].

Postoperative complications were defined differently in included trials, most often as any kind of prosthetic or biological complications. A systematic review indicated

that biological and technical complications following dental implant therapy should be better specified [164].

This study included only RCTs included in available Cochrane SRs, which may be regarded as a limitation. It is acknowledged that there could be relevant RCTs that were not included in the analyzed SRs. However, considering the comprehensive search and robust methodology of Cochrane SRs, as well as the wide period during which the trials analyzed in our study were published (from 1995 to 2020), it is highly likely that these reviews covered clinical questions of highest priority and included all relevant clinical trials concerning implant prosthodontics. Also, we made arbitrary decisions when designing this study, because there are no methodological guidelines for conducting analyses of outcomes. Most importantly, increasing the sample of studies in this work is unlikely to change the results. We have already proven on this sample that there is a very high heterogeneity of methodological approaches in the studied field.

This study can be helpful for the next step of developing a COS for dental implant therapy. The development and application of an agreed COS have been spearheaded by the COMET (Core Outcome Measures in Effectiveness Trials) Initiative, which was launched in 2010 [19]. The COS is mainly developed by relevant experts in a particular field of medicine, but it also involves the public to ensure that those outcomes are relevant and important to patients. If the findings are to influence policy and practice, then the chosen outcomes need to be relevant and important to key stakeholders, including patients and the public, health care professionals, and others making decisions about health care [165].

When searching the literature, we found no similar research that took a methodological look at the diversity of outcomes in research on dental implant prosthetics.

The COS in the field of implant prosthodontics has not been proposed yet. Therefore, our results have the potential to contribute to the development of COS and improve the use of relevant outcomes in future trials on implant prosthodontic therapy. The use of COS would lead to a standardized use of outcomes, enable future synthesis and comparison of the obtained results within different studies and preserve resources for the improvement of future research.

The creation and use of COS in studies leads to a reduction in the number of studies that are considered research waste, which is defined as studies that are not necessary, and have poor design, conduct or reporting. Research waste hinders or prevents the synthesis and usability of study results [166, 167].

Conclusions

Clinical trials in implant prosthodontics used different outcomes, different definitions of outcomes and used different times to monitor them. Standardization of outcomes is necessary to allow comparability and evidence synthesis about the effectiveness of implant prosthodontic therapy. Future research should include defining a core outcome set for implant prosthodontic therapy.

List of abbreviations

CDP	Complete dental prostheses
COHG	Cochrane Oral Health Group
COMET	Core Outcome Measures in Effectiveness Trials Initiative
COS	Core outcome set
FCPD	Fixed complete dental prostheses
FPD	Fixed partial dentures
NSAID	Non-steroidal anti-inflammatory drugs
PES	Pink Aesthetic Scores
RCT	Randomized controlled trial
STROBE	STrengthening the Reporting of OBServational studies in Epidemiology
WES	White Esthetic Score

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12903-023-03658-9>.

Supplementary Material 1: Study protocol
Supplementary Material 2: List of included systematic reviews and randomized controlled trials
Supplementary Material 3: Countries in which randomized controlled trials were conducted
Supplementary Material 4: Raw data extractions

Acknowledgements

We are thankful to professor Marco Esposito for providing us with the full text of the study we needed for our analyses. This study was part of the graduation thesis of Ante Vardić at the University of Split School of Medicine Dental Medicine Study. The graduation thesis was written in Croatian. Preliminary results of this study were presented as an oral presentation at the Congress organised by the Croatian Society of Periodontology in Rovinj, Croatia on November 25th 2022.

Authors' contributions

Study design: AV, LP, TPP. Data collection and analyses: AV, TG, JV, EK, TPP. Writing the first draft of the manuscript: AV, LP, TPP. Critical revision of the manuscript: AV, LP, TG, JV, EK, TPP. Approval of the final version of the manuscript: All authors read and approved the final manuscript.

Funding

The authors received no funding for this work.

Data Availability

Raw data generated by extractions from the included trials and all relevant data concerning SRs or RCTs included in our analyses are available in Supplementary file 4.

Declarations

Ethics approval and consent to participate

Not applicable – this was analysis of articles published in scholarly journals.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Study of Dental Medicine, University of Split School of Medicine, Split, Croatia

²Center for Evidence-Based Medicine and Health Care, Catholic University of Croatia, Zagreb, Croatia

³Department of Prosthodontics, Study of Dental Medicine, University of Split School of Medicine, Split, Croatia

⁴Department of Fixed Prosthodontics, University of Zagreb School of Dental Medicine, Zagreb, Croatia

Received: 3 January 2023 / Accepted: 12 November 2023

Published online: 22 November 2023

References

- Alghamdi HS, Jansen JA. The development and future of dental implants. *Dent Mater J*. 2020;39(2):167–72.
- Hartlev J, Kohberg P, Ahlmann S, Andersen NT, Schou S, Isidor F. Patient satisfaction and esthetic outcome after immediate placement and provisionalization of single-tooth implants involving a definitive individual abutment. *Clin Oral Implants Res*. 2014;25(11):1245–50.
- Jofre J, Castiglioni X, Lobos CA. Influence of minimally invasive implant-retained overdenture on patients' quality of life: a randomized clinical trial. *Clin Oral Implants Res*. 2013;24(10):1173–7.
- Shemtov-Yona K, Rittel D. An overview of the Mechanical Integrity of Dental Implants. *Biomed Res Int*. 2015;2015:547384.
- Esposito M, Hirsch JM, Lekholm U, Thomsen P. Biological factors contributing to failures of osseointegrated oral implants. (I). Success criteria and epidemiology. *Eur J Oral Sci*. 1998;106(1):527–51.
- Tonetti MS, Schmid J. Pathogenesis of implant failures. *Periodontol* 2000. 1994;4:127–38.
- Manor Y, Oubaid S, Mardinger O, Chaushu G, Nissan J. Characteristics of early versus late implant failure: a retrospective study. *J Oral Maxillofac Surg*. 2009;67(12):2649–52.
- Sakka S, Baroudi K, Nassani MZ. Factors associated with early and late failure of dental implants. *J Investig Clin Dent*. 2012;3(4):258–61.
- Cosola S, Toti P, Babetto E, Covani U, Peñarrocha-Diogo M, Peñarrocha-Oltra D. In-vitro fatigue and fracture performance of three different ferulized implant connections used in fixed prosthesis. *J Dent Sci*. 2021;16(1):397–403.
- Clark D, Levin L. Dental implant management and maintenance: how to improve long-term implant success? *Quintessence Int*. 2016;47(5):417–23.
- Shibuya Y, Takata N, Takeuchi J, Tsuji K, Ishida S, Kobayashi M, Suzuki H, Hasegawa T, Kamae I, Komori T. Analysis of the 619 Brånemark System TiUnite implants: a retrospective study. *Kobe J Med Sci*. 2012;58(1):E19–28.
- Albrektsson T, Zarb G, Worthington P, Eriksson AR. The long-term efficacy of currently used dental implants: a review and proposed criteria of success. *Int J Oral Maxillofac Implants*. 1986;1(1):11–25.
- Misch CE, Perel ML, Wang HL, Sammartino G, Galindo-Moreno P, Trisi P, Steigmann M, Rebaudi A, Palti A, Pikos MA, Schwartz-Arad D, Choukroun J, Gutierrez-Perez JL, Marenzi G, Valavanis DK. Implant success, survival, and failure: the International Congress of Oral Implantologists (ICOI) Pisa Consensus Conference. *Implant Dent*. 2008;17(1):5–15.
- Annibaldi S, Bignozzi I, La Monaca G, Cristalli MP. Usefulness of the aesthetic result as a success criterion for implant therapy: a review. *Clin Implant Dent Relat Res*. 2012;14(1):3–40.
- Buser D, Mericske-Stern R, Bernard JP, Behneke A, Behneke N, Hirt HP, Belser UC, Lang NP. Long-term evaluation of non-submerged ITI implants. Part 1: 8-year life table analysis of a prospective multi-center study with 2359 implants. *Clin Oral Implants Res*. 1997;8(3):161–72.
- Lindeboom JA, Frenken JW, Dubois L, Frank M, Abbink I, Kroon FH. Immediate loading versus immediate provisionalization of maxillary single-tooth replacements: a prospective randomized study with BioComp implants. *J Oral Maxillofac Surg*. 2006;64(6):936–42.
- Fürhauser R, Florescu D, Benesch T, Haas R, Mailath G, Watzek G. Evaluation of soft tissue around single-tooth implant crowns: the pink esthetic score. *Clin Oral Implants Res*. 2005;16(6):639–44.

18. Meijer HJ, Stellingsma K, Meijndert L, Raghoobar GM. A new index for rating aesthetics of implant-supported single crowns and adjacent soft tissues—the Implant Crown Aesthetic Index. *Clin Oral Implants Res.* 2005;16(6):645–9.
19. Belser UC, Grütter L, Vailati F, Bornstein MM, Weber HP, Buser D. Outcome evaluation of early placed maxillary anterior single-tooth implants using objective esthetic criteria: a cross-sectional, retrospective study in 45 patients with a 2- to 4-year follow-up using pink and white esthetic scores. *J Periodontol.* 2009;80(1):140–51.
20. COMET (Core Outcome Measures in Effectiveness Trials) Initiative. <http://www.comet-initiative.org/>
21. Jambi S, Walsh T, Sandler J, Benson PE, Skeggs RM, O'Brien KD. Reinforcement of anchorage during orthodontic brace treatment with implants or other surgical methods. *Cochrane Database Syst Rev.* 2014;2014(8):CD005098.
22. El-Rabbany M, Duchnay M, Raziee HR, Zych M, Tenenbaum H, Shah PS, Azarpazhooh A. Interventions for preventing osteoradionecrosis of the jaws in adults receiving head and neck radiotherapy. *Cochrane Database Syst Rev.* 2019;2019(11):CD011559.
23. Esposito M, Worthington HV. Interventions for replacing missing teeth: dental implants in zygomatic bone for the rehabilitation of the severely deficient edentulous maxilla. *Cochrane Database Syst Rev.* 2013;2013(9):CD004151.
24. Esposito M, Grusovin MG, Worthington HV. Interventions for replacing missing teeth: antibiotics at dental implant placement to prevent Complications. *Cochrane Database of Systematic Reviews* 2013, Issue 7. Art. No.: CD004152.
25. Payne AGT, Alsabeeha NHM, Atieh MA, Esposito M, Ma S, Anas El-Wegoud M. Interventions for replacing missing teeth: attachment systems for implant overdentures in edentulous jaws. *Cochrane Database of Systematic Reviews.* 2018; Issue 10: ArtNoCD008001.
26. Esposito M, Felice P, Worthington HV. Interventions for replacing missing teeth: augmentation procedures of the maxillary sinus. *Cochrane Database of Systematic Reviews.* Issue 2014;5:CD008397. Art. No.
27. Esposito M, Grusovin MG, Polyzos IP, Felice P, Worthington HV. Interventions for replacing missing teeth: dental implants in fresh extraction sockets (immediate, immediate-delayed and delayed implants). *Cochrane Database of Systematic Reviews.* 2010; Issue 9: ArtNoCD005968.
28. Esposito M, Grusovin MG, Maghairieh H, Worthington HV. Interventions for replacing missing teeth: different times for loading dental implants. *Cochrane Database of Systematic Reviews* 2013, Issue 3. Art. No.: CD003878.
29. Esposito M, Grusovin MG, Felice P, Karatzopoulos G, Worthington HV, Coulthard P. Interventions for replacing missing teeth: horizontal and vertical bone augmentation techniques for dental implant treatment. *Cochrane Database of Systematic Reviews* 2009, Issue 4. Art. No.: CD003607.
30. Esposito M, Worthington HV. Interventions for replacing missing teeth: hyperbaric oxygen therapy for irradiated patients who require dental implants. *Cochrane Database of Systematic Reviews* 2013, Issue 9. Art. No.: CD003603.
31. Grusovin MG, Coulthard P, Worthington HV, George P, Esposito M. Interventions for replacing missing teeth: maintaining and recovering soft tissue health around dental implants. *Cochrane Database of Systematic Reviews* 2010, Issue 8. Art. No.: CD003069.
32. Esposito M, Maghairieh H, Grusovin MG, Ziounas I, Worthington HV. Interventions for replacing missing teeth: management of soft tissues for dental implants. *Cochrane Database of Systematic Reviews* 2012, Issue 2. Art. No.: CD006697.
33. Esposito M, Grusovin MG, Worthington HV. Interventions for replacing missing teeth: treatment of peri-implantitis. *Cochrane Database of Systematic Reviews.* Issue 2012;1:CD004970. Art. No.
34. Atieh MA, Alsabeeha NHM, Payne AGT, Ali S, Faggion CM Jr, Esposito M. Interventions for replacing missing teeth: alveolar ridge preservation techniques for dental implant site development. *Cochrane Database of Systematic Reviews* 2021, Issue 4. Art. No.: CD010176.
35. Raghoobar GM, Schortinghuis J, Liem RS, Ruben JL, van der Wal JE, Vissink A. Does platelet-rich plasma promote remodeling of autologous bone grafts used for augmentation of the maxillary sinus floor? *Clin Oral Implants Res.* 2005;16(3):349–56.
36. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Syst Reviews.* 2021;10(1):1–1.
37. Brkovic BM, Prasad HS, Rohrer MD, Konandreas G, Agrogiannis G, Antunovic D, Sándor GK. Beta-tricalcium phosphate/type I collagen cones with or without a barrier membrane in human extraction socket healing: clinical, histologic, histomorphometric, and immunohistochemical evaluation. *Clin Oral Investig.* 2012;16(2):581–90.
38. Cha JK, Song YW, Park SH, Jung RE, Jung UW, Thoma DS. Alveolar ridge preservation in the posterior maxilla reduces vertical dimensional change: a randomized controlled clinical trial. *Clin Oral Implants Res.* 2019;30(6):515–23.
39. Fernandes PG, Novaes AB Jr, de Queiroz AC, de Souza SL, Taba M Jr, Palioto DB, Grisi MF. Ridge preservation with acellular dermal matrix and anorganic bone matrix cell-binding peptide P-15 after tooth extraction in humans. *J Periodontol.* 2011;82(1):72–9.
40. Festa VM, Addabbo F, Laino L, Femiano F, Rullo R. Porcine-derived xenograft combined with a soft cortical membrane versus extraction alone for implant site development: a clinical study in humans. *Clin Implant Dent Relat Res.* 2013;15(5):707–13.
41. Hoang TN, Mealey BL. Histologic comparison of healing after ridge preservation using human demineralized bone matrix putty with one versus two different-sized bone particles. *J Periodontol.* 2012;83(2):174–81.
42. Iorio-Siciliano V, Blasi A, Nicolò M, Iorio-Siciliano A, Riccitiello F, Ramaglia L. Clinical Outcomes of Socket Preservation Using Bovine-Derived Xenograft Collagen and Collagen Membrane post-tooth extraction: a 6-Month Randomized Controlled Clinical Trial. *Int J Periodontics Restorative Dent* 2017 Sep/Oct;37(5):e290–6.
43. Cepa S, Koller B, Spies BC, Stampf S, Kohal RJ. Implant-retained prostheses: ball vs. conus attachments - a randomized controlled clinical trial. *Clin Oral Implants Res.* 2017;28(2):177–85.
44. Kappel S, Giannakopoulos NN, Eberhard L, Rammelsberg P, Eiffler C. Immediate Loading of Dental implants in Edentulous mandibles by Use of Locator® attachments or Dolder® bars: two-year results from a prospective Randomized Clinical Study. *Clin Implant Dent Relat Res.* 2016;18(4):752–61.
45. Naert I, Gizani S, Vuylsteke M, Van Steenberghe D. A 5-year prospective randomized clinical trial on the influence of splinted and unsplinted oral implants retaining a mandibular overdenture: prosthetic aspects and patient satisfaction. *J Oral Rehabil.* 1999;26(3):195–202.
46. Tawse-Smith A, Payne AG, Kumara R, Thomson WM. Early loading of unsplinted implants supporting mandibular overdentures using a one-stage operative procedure with two different implant systems: a 2-year report. *Clin Implant Dent Relat Res.* 2002;4(1):33–42.
47. Abu-Ta'a M, Quirynen M, Teughels W, van Steenberghe D. Asepsis during periodontal Surgery involving oral implants and the usefulness of peri-operative antibiotics: a prospective, randomized, controlled clinical trial. *J Clin Periodontol.* 2008;35(1):58–63.
48. Caiazzo A, Casavecchia P, Barone A, Brugnamì F. A pilot study to determine the effectiveness of different Amoxicillin regimens in implant Surgery. *J Oral Implantol.* 2011;37(6):691–6.
49. Esposito M, Cannizzaro G, Bozzoli P, Consolo U, Felice P, Ferri V, Landriani S, Leone M, Magliano A, Pellitteri G, Todisco M, Torchio C. Efficacy of prophylactic antibiotics for dental implants: a multicentre placebo-controlled randomised clinical trial. *Eur J Oral Implantol.* 2008;1(1):23–31.
50. Esposito M, Cannizzaro G, Bozzoli P, Checchi L, Ferri V, Landriani S, Leone M, Todisco M, Torchio C, Testori T, Galli F, Felice P. Effectiveness of prophylactic antibiotics at placement of dental implants: a pragmatic multi-centre placebo-controlled randomised clinical trial. *Eur J Oral Implantol.* 2010;3(2):135–43.
51. Nolan R, Kemmoona M, Polyzos I, Claffey N. The influence of prophylactic antibiotic administration on post-operative morbidity in dental implant Surgery. A prospective double blind randomized controlled clinical trial. *Clin Oral Implants Res.* 2014;25(2):252–9.
52. Cannizzaro G, Felice P, Leone M, Viola P, Esposito M. Early loading of implants in the atrophic posterior maxilla: lateral sinus lift with autogenous bone and Bio-oss versus crestal mini sinus lift and 8-mm hydroxyapatite-coated implants. A randomised controlled clinical trial. *Eur J Oral Implantol.* 2009;2(1):25–38.
53. Checchi L, Felice P, Antonini ES, Cosci F, Pellegrino G, Esposito M. Crestal sinus lift for implant rehabilitation: a randomised clinical trial comparing the Cosci and the summers techniques. A preliminary report on Complications and patient preference. *Eur J Oral Implantol.* 2010;3(3):221–32.
54. Crespi R, Capparè P, Gherlone E. Sinus floor elevation by osteotome: hand mallet versus electric mallet. A prospective clinical study. *Int J Oral Maxillofac Implants.* 2012;27(5):1144–50.
55. Esposito M, Cannizzaro G, Soardi E, Pistilli R, Piattelli M, Corvino V, Felice P. Posterior atrophic jaws rehabilitated with prostheses supported by 6 mm-long, 4 mm-wide implants or by longer implants in augmented bone. Preliminary results from a pilot randomised controlled trial. *Eur J Oral Implantol.* 2012;5(1):19–33.

56. Felice P, Checchi V, Pistilli R, Scarano A, Pellegrino G, Esposito M. Bone augmentation versus 5-mm dental implants in posterior atrophic jaws. Four-month post-loading results from a randomised controlled clinical trial. *Eur J Oral Implantol*. 2009;2(4):267–81.
57. Felice P, Scarano A, Pistilli R, Checchi L, Piattelli M, Pellegrino G, Esposito M. A comparison of two techniques to augment maxillary sinuses using the lateral window approach: rigid synthetic resorbable barriers versus anorganic bovine bone. Five-month post-loading clinical and histological results of a pilot randomised controlled clinical trial. *Eur J Oral Implantol*. 2009;2(4):293–306.
58. Felice P, Soardi E, Pellegrino G, Pistilli R, Marchetti C, Gessaroli M, Esposito M. Treatment of the atrophic edentulous maxilla: short implants versus bone augmentation for placing longer implants. Five-month post-loading results of a pilot randomised controlled trial. *Eur J Oral Implantol*. 2011;4(3):191–202.
59. Felice P, Pistilli R, Piattelli M, Soardi E, Corvino V, Esposito M. Posterior atrophic jaws rehabilitated with prostheses supported by 5 x 5 mm implants with a novel nanostructured calcium-incorporated titanium surface or by longer implants in augmented bone. Preliminary results from a randomised controlled trial. *Eur J Oral Implantol*. 2012;5(2):149–61.
60. Felice P, Pistilli R, Piattelli M, Soardi E, Pellegrino G, Corvino V, Esposito M. 1-stage versus 2-stage lateral maxillary sinus lift procedures: 4-month post-loading results of a multicenter randomised controlled trial. *Eur J Oral Implantol*. 2013;6(2):153–65.
61. Lindgren C, Mordenfeld A, Hallman M. A prospective 1-year clinical and radiographic study of implants placed after maxillary sinus floor augmentation with synthetic biphasic calcium phosphate or deproteinized bovine bone. *Clin Implant Dent Relat Res*. 2012;14(1):41–50.
62. Merli M, Moscatelli M, Mariotti G, Rotundo R, Nieri M. Autogenous bone versus deproteinised bovine bone matrix in 1-stage lateral sinus floor elevation in the severely atrophied maxilla: a randomised controlled trial. *Eur J Oral Implantol*. 2013;6(1):27–37.
63. Si MS, Zhuang LF, Gu YX, Mo JJ, Qiao SC, Lai HC. Osteotome sinus floor elevation with or without grafting: a 3-year randomized controlled clinical trial. *J Clin Periodontol*. 2013;40(4):396–403.
64. Torres J, Tamimi F, Martinez PP, Alkhrasat MH, Linares R, Hernández G, Torres-Macho J, López-Cabarcos E. Effect of platelet-rich plasma on sinus lifting: a randomized-controlled clinical trial. *J Clin Periodontol*. 2009;36(8):677–87.
65. Torres García-Denche J, Wu X, Martinez PP, Eimar H, Ikbal DJ, Hernández G, López-Cabarcos E, Fernandez-Tresguerres I, Tamimi F. Membranes over the lateral window in sinus augmentation procedures: a two-arm and split-mouth randomized clinical trials. *J Clin Periodontol*. 2013;40(11):1043–51.
66. Wannfors K, Johansson B, Hallman M, Strandkvist T. A prospective randomized study of 1- and 2-stage sinus inlay bone grafts: 1-year follow-up. *Int J Oral Maxillofac Implants*. 2000 Sep-Oct;15(5):625–32.
67. Chen ST, Darby IB, Adams GG, Reynolds EC. A prospective clinical study of bone augmentation techniques at immediate implants. *Clin Oral Implants Res*. 2005;16(2):176–84.
68. Chen ST, Darby IB, Reynolds EC. A prospective clinical study of non-submerged immediate implants: clinical outcomes and esthetic results. *Clin Oral Implants Res*. 2007;18(5):552–62.
69. Lindeboom JA, Tjiook Y, Kroon FH. Immediate placement of implants in periapical infected sites: a prospective randomized study in 50 patients. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2006;101(6):705–10.
70. Assad AS, Hassan SA, Shawky YM, Badawy MM. Clinical and radiographic evaluation of implant-retained mandibular overdentures with immediate loading. *Implant Dent*. 2007;16(2):212–23.
71. Cannizzaro G, Leone M. Restoration of partially edentulous patients using dental implants with a microtextured surface: a prospective comparison of delayed and immediate full occlusal loading. *Int J Oral Maxillofac Implants*. 2003 Jul-Aug;18(4):512–22.
72. Cannizzaro G, Leone M, Esposito M. Immediate versus early loading of two implants placed with a flapless technique supporting mandibular bar-retained overdentures: a single-blinded, randomised controlled clinical trial. *Eur J Oral Implantol*. 2008;1(1):33–43.
73. Cannizzaro G, Torchio C, Leone M, Esposito M. Immediate versus early loading of flapless-placed implants supporting maxillary full-arch prostheses: a randomised controlled clinical trial. *Eur J Oral Implantol*. 2008;9(Suppl 1):127–39.
74. Cannizzaro G, Leone M, Torchio C, Viola P, Esposito M. Immediate versus early loading of 7-mm-long flapless-placed single implants: a split-mouth randomised controlled clinical trial. *Eur J Oral Implantol*. 2008;1(4):277–92.
75. Cannizzaro G, Torchio C, Felice P, Leone M, Esposito M. Immediate occlusal versus non-occlusal loading of single zirconia implants. A multicentre pragmatic randomised clinical trial. *Eur J Oral Implantol*. 2010;3(2):111–20.
76. Chiapasco M, Abati S, Romeo E, Vogel G. Implant-retained mandibular overdentures with Brånemark System MKII implants: a prospective comparative study between delayed and immediate loading. *Int J Oral Maxillofac Implants*. 2001 Jul-Aug;16(4):537–46.
77. Crespi R, Capparé P, Gherlone E, Romanos GE. Immediate versus delayed loading of dental implants placed in fresh extraction sockets in the maxillary esthetic zone: a clinical comparative study. *Int J Oral Maxillofac Implants*. 2008 Jul-Aug;23(4):753–8.
78. Donati M, La Scala V, Billi M, Di Dino B, Torrisi P, Berglundh T. Immediate functional loading of implants in single tooth replacement: a prospective clinical multicenter study. *Clin Oral Implants Res*. 2008;19(8):740–8.
79. Fischer K, Stenberg T. Early loading of ITI implants supporting a maxillary full-arch prosthesis: 1-year data of a prospective, randomized study. *Int J Oral Maxillofac Implants*. 2004 May-Jun;19(3):374–81.
80. Güncü MB, Aslan Y, Tümer C, Güncü GN, Uysal S. In-patient comparison of immediate and conventional loaded implants in mandibular molar sites within 12 months. *Clin Oral Implants Res*. 2008;19(4):335–41.
81. Hall JA, Payne AG, Purton DG, Torr B. A randomized controlled clinical trial of conventional and immediately loaded tapered implants with screw-retained crowns. *Int J Prosthodont*. 2006 Jan-Feb;19(1):17–9.
82. Merli M, Merli A, Bernardelli F, Lombardini F, Esposito M. Immediate versus early non-occlusal loading of dental implants placed flapless in partially edentulous patients. One-year results from a randomised controlled trial. *Eur J Oral Implantol*. 2008;1(3):207–20.
83. Oh TJ, Shotwell JL, Billy EJ, Wang HL. Effect of flapless implant surgery on soft tissue profile: a randomized controlled clinical trial. *J Periodontol*. 2006;77(5):874–82.
84. Payne AG, Tawse-Smith A, Duncan WD, Kumara R. Conventional and early loading of unsplinted ITI implants supporting mandibular overdentures. *Clin Oral Implants Res*. 2002;13(6):603–9.
85. Schincaglia GP, Marzola R, Giovanni GF, Chiara CS, Scotti R. Replacement of mandibular molars with single-unit restorations supported by wide-body implants: immediate versus delayed loading. A randomized controlled study. *Int J Oral Maxillofac Implants*. 2008 May-Jun;23(3):474–80.
86. Testori T, Galli F, Capelli M, Zuffetti F, Esposito M. Immediate nonocclusal versus early loading of dental implants in partially edentulous patients: 1-year results from a multicenter, randomized controlled clinical trial. *Int J Oral Maxillofac Implants*. 2007 Sep-Oct;22(5):815–22.
87. Turkyilmaz I, Tumer C. Early versus late loading of unsplinted TiUnite surface implants supporting mandibular overdentures: a 2-year report from a prospective study. *J Oral Rehabil*. 2007;34(10):773–80.
88. Zöllner A, Ganeles J, Korostoff J, Guerra F, Krafft T, Brägger U. Immediate and early non-occlusal loading of Straumann implants with a chemically modified surface (SLActive) in the posterior mandible and maxilla: interim results from a prospective multicenter randomized-controlled study. *Clin Oral Implants Res*. 2008;19(5):442–50.
89. Bianchi A, Felice P, Lizio G, Marchetti C. Alveolar distraction osteogenesis versus inlay bone grafting in posterior mandibular atrophy: a prospective study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2008;105(3):282–92.
90. Chiapasco M, Romeo E, Casentini P, Rimondini L. Alveolar distraction osteogenesis vs. vertical guided bone regeneration for the correction of vertically deficient edentulous ridges: a 1-3-year prospective study on humans. *Clin Oral Implants Res*. 2004;15(1):82–95.
91. Chiapasco M, Zaniboni M, Rimondini L. Autogenous onlay bone grafts vs. alveolar distraction osteogenesis for the correction of vertically deficient edentulous ridges: a 2-4-year prospective study on humans. *Clin Oral Implants Res*. 2007;18(4):432–40.
92. Felice P, Marchetti C, Piattelli A, Pellegrino G, Checchi V, Worthington H, Esposito M. Vertical ridge augmentation of the atrophic posterior mandible with interpositional block grafts: bone from the iliac crest versus bovine anorganic bone. *Eur J Oral Implantol*. 2008;1(3):183–98.
93. Felice P, Cannizzaro G, Checchi V, Marchetti C, Pellegrino G, Censi P, Esposito M. Vertical bone augmentation versus 7-mm-long implants in posterior atrophic mandibles. Results of a randomised controlled clinical trial of up to 4 months after loading. *Eur J Oral Implantol*. 2009;2(1):7–20.
94. Felice P, Pistilli R, Lizio G, Pellegrino G, Nisii A, Marchetti C. Inlay versus onlay iliac bone grafting in atrophic posterior mandible: a prospective controlled clinical trial for the comparison of two techniques. *Clin Implant Dent Relat Res*. 2009;11(Suppl 1):e69–82.

95. Fontana F, Santoro F, Maiorana C, Iezzi G, Piattelli A, Simion M. Clinical and histologic evaluation of allogeneic bone matrix versus autogenous bone chips associated with titanium-reinforced e-PTFE membrane for vertical ridge augmentation: a prospective pilot study. *Int J Oral Maxillofac Implants*. 2008 Nov-Dec;23(6):1003–12.
96. Merli M, Migani M, Esposito M. Vertical ridge augmentation with autogenous bone grafts: resorbable barriers supported by osteosynthesis plates versus titanium-reinforced barriers. A preliminary report of a blinded, randomized controlled clinical trial. *Int J Oral Maxillofac Implants*. 2007 May-Jun;22(3):373–82.
97. Raghoobar GM, Liem RS, Bos RR, van der Wal JE, Vissink A. Resorbable screws for fixation of autologous bone grafts. *Clin Oral Implants Res*. 2006;17(3):288–93.
98. Schortinghuis J, Bronckers AL, Gravendeel J, Stegenga B, Raghoobar GM. The effect of ultrasound on osteogenesis in the vertically distracted edentulous mandible: a double-blind trial. *Int J Oral Maxillofac Surg*. 2008;37(11):1014–21.
99. Stellingsma K, Bouma J, Stegenga B, Meijer HJ, Raghoobar GM. Satisfaction and psychosocial aspects of patients with an extremely resorbed mandible treated with implant-retained overdentures. A prospective, comparative study. *Clin Oral Implants Res*. 2003;14(2):166–72.
100. de Araújo Nobre M, Cintra N, Maló P. Peri-implant maintenance of immediate function implants: a pilot study comparing hyaluronic acid and chlorhexidine. *Int J Dent Hyg*. 2007;5(2):87–94.
101. Arnabat-Domínguez J, España-Tost AJ, Berini-Aytés L, Gay-Escoda C. Erbium:YAG laser application in the second phase of implant Surgery: a pilot study in 20 patients. *Int J Oral Maxillofac Implants*. 2003 Jan-Feb;18(1):104–12.
102. Cannizzaro G, Felice P, Leone M, Checchi V, Esposito M. Flapless versus open flap implant Surgery in partially edentulous patients subjected to immediate loading: 1-year results from a split-mouth randomised controlled trial. *Eur J Oral Implantol*. 2011;4(3):177–88.
103. Heydenrijk K, Raghoobar GM, Batenburg RH, Stegenga B. A comparison of labial and crestal incisions for the 1-stage placement of IMZ implants: a pilot study. *J Oral Maxillofac Surg*. 2000;58(10):1119–23. discussion 1123–4.
104. Lindeboom JA, van Wijk AJ. A comparison of two implant techniques on patient-based outcome measures: a report of flapless vs. conventional flapped implant placement. *Clin Oral Implants Res*. 2010;21(4):366–70.
105. Lorenzo R, García V, Orsini M, Martin C, Sanz M. Clinical efficacy of a xenogenic collagen matrix in augmenting keratinized mucosa around implants: a randomized controlled prospective clinical trial. *Clin Oral Implants Res*. 2012;23(3):316–24.
106. Wiesner G, Esposito M, Worthington H, Schlee M. Connective tissue grafts for thickening peri-implant tissues at implant placement. One-year results from an explanatory split-mouth randomised controlled clinical trial. *Eur J Oral Implantol*. 2010;3(1):27–35.
107. Büchter A, Meyer U, Kruse-Lösler B, Joos U, Kleinheinz J. Sustained release of doxycycline for the treatment of peri-implantitis: randomised controlled trial. *Br J Oral Maxillofac Surg*. 2004;42(5):439–44.
108. Karring ES, Stavropoulos A, Ellegaard B, Karring T. Treatment of peri-implantitis by the Vector system. *Clin Oral Implants Res*. 2005;16(3):288–93.
109. Renvert S, Lindahl C, Roos Jansäker AM, Persson GR. Treatment of peri-implantitis using an Er:YAG laser or an air-abrasive device: a randomized clinical trial. *J Clin Periodontol*. 2011;38(1):65–73.
110. Romeo E, Ghisolfi M, Murgolo N, Chiapasco M, Lops D, Vogel G. Therapy of peri-implantitis with resective Surgery. A 3-year clinical trial on rough screw-shaped oral implants. Part I: clinical outcome. *Clin Oral Implants Res*. 2005;16(1):9–18.
111. Schwarz F, Sculean A, Rothamel D, Schwenzler K, Georg T, Becker J. Clinical evaluation of an Er:YAG laser for nonsurgical treatment of peri-implantitis: a pilot study. *Clin Oral Implants Res*. 2005;16(1):44–52.
112. Schwarz F, Bieling K, Bonsmann M, Latz T, Becker J. Nonsurgical treatment of moderate and advanced periimplantitis lesions: a controlled clinical study. *Clin Oral Investig*. 2006;10(4):279–88.
113. Schwarz F, Bieling K, Latz T, Nuesry E, Becker J. Healing of intrabony peri-implantitis defects following application of a nanocrystalline hydroxyapatite (Ostim) or a bovine-derived xenograft (Bio-Oss) in combination with a collagen membrane (Bio-Gide). A case series. *J Clin Periodontol*. 2006;33(7):491–9.
114. Schwarz F, Sahn N, Iglhaut G, Becker J. Impact of the method of surface debridement and decontamination on the clinical outcome following combined surgical therapy of peri-implantitis: a randomized controlled clinical study. *J Clin Periodontol*. 2011;38(3):276–84.
115. Tang Z, Cao C, Sha Y, Lin Y, Wang X. Effects of non-surgical treatment modalities on peri-implantitis. *Zhonghua Kou Qiang Yi Xue Za Zhi*. 2002;37(3):173–5.
116. Barone A, Orlando B, Cingano L, Marconcini S, Derchi G, Covani U. A randomized clinical trial to evaluate and compare implants placed in augmented versus non-augmented extraction sockets: 3-year results. *J Periodontol*. 2012;83(7):836–46.
117. Block MS, Mercante DE, Lirette D, Mohamed W, Ryser M, Castellon P. Prospective evaluation of immediate and delayed provisional single tooth restorations. *J Oral Maxillofac Surg*. 2009;67(11 Suppl):89–107.
118. De Rouck T, Collys K, Wyn I, Cosyn J. Instant provisionalization of immediate single-tooth implants is essential to optimize esthetic treatment outcome. *Clin Oral Implants Res*. 2009;20(6):566–70.
119. Enkling N, Albrecht D, Bayer S, Stark H, Mericske-Stern R. [Immediate loading of interforaminal implants using a chairside fabricated bur (oral presentation)]. *Clin Oral Implants Res*. 2010;21:1013.
120. Horwitz J, Machtei EE, Zuabi O, Peled M. Amine fluoride/stannous fluoride and chlorhexidine mouthwashes as adjuncts to single-stage dental implants: a comparative study. *J Periodontol*. 2005;76(3):334–40.
121. Meijndert L, Meijer HJ, Stellingsma K, Stegenga B, Raghoobar GM. Evaluation of aesthetics of implant-supported single-tooth replacements using different bone augmentation procedures: a prospective randomized clinical study. *Clin Oral Implants Res*. 2007;18(6):715–9.
122. Rickert D, Vissink A, Slater JJ, Meijer HJ, Raghoobar GM. Comparison between conventional and piezoelectric surgical tools for maxillary sinus floor elevation. A randomized controlled clinical trial. *Clin Implant Dent Relat Res*. 2013;15(2):297–302.
123. Schoen PJ, Raghoobar GM, Bouma J, Reintsema H, Vissink A, Sterk W, Roodenburg JL. Rehabilitation of oral function in Head and Neck cancer patients after radiotherapy with implant-retained dentures: effects of hyperbaric oxygen therapy. *Oral Oncol*. 2007;43(4):379–88.
124. Patel K, Mardas N, Donos N. Radiographic and clinical outcomes of implants placed in ridge preserved sites: a 12-month post-loading follow-up. *Clin Oral Implants Res*. 2013;24(6):599–605.
125. Cochran DL, Buser D, ten Bruggenkate CM, Weingart D, Taylor TM, Bernard JP, Peters F, Simpson JP. The use of reduced healing times on ITI implants with a sandblasted and acid-etched (SLA) surface: early results from clinical trials on ITI SLA implants. *Clin Oral Implants Res*. 2002;13(2):144–53.
126. Hallman M, Sennerby L, Lundgren S. A clinical and histologic evaluation of implant integration in the posterior maxilla after sinus floor augmentation with autogenous bone, bovine hydroxyapatite, or a 20:80 mixture. *Int J Oral Maxillofac Implants*. 2002 Sep-Oct;17(5):635–43.
127. Palatella P, Torsello F, Cordaro L. Two-year prospective clinical comparison of immediate replacement vs. immediate restoration of single tooth in the esthetic zone. *Clin Oral Implants Res*. 2008;19(11):1148–53.
128. Schropp L, Kostopoulos L, Wenzel A. Bone healing following immediate versus delayed placement of titanium implants into extraction sockets: a prospective clinical study. *Int J Oral Maxillofac Implants*. 2003 Mar-Apr;18(2):189–99.
129. Cristache CM, Muntianu LA, Burlibasa M, Didilescu AC. Five-year clinical trial using three attachment systems for implant overdentures. *Clin Oral Implants Res*. 2014;25(2):e171–8.
130. Cune M, Burgers M, van Kampen F, de Putter C, van der Bilt A. Mandibular overdentures retained by two implants: 10-year results from a crossover clinical trial comparing ball-socket and bar-clip attachments. *Int J Prosthodont*. 2010 Jul-Aug;23(4):310–7.
131. Walton JN, MacEntee MI, Glick N. One-year prosthetic outcomes with implant overdentures: a randomized clinical trial. *Int J Oral Maxillofac Implants*. 2002 May-Jun;17(3):391–8.
132. Anitua E, Aguirre JJ, Gorosabel A, Barrio P, Errazquin JM, Román P, Pla R, Carrete J, de Petro J, Orive G. A multicentre placebo-controlled randomised clinical trial of antibiotic prophylaxis for placement of single dental implants. *Eur J Oral Implantol*. 2009;2(4):283–92.
133. Meloni SM, De Riu G, Pisano M, De Riu N, Tullio A. Immediate versus delayed loading of single mandibular molars. One-year results from a randomised controlled trial. *Eur J Oral Implantol*. 2012;5(4):345–53.
134. Renvert S, Samuelsson E, Lindahl C, Persson GR. Mechanical non-surgical treatment of peri-implantitis: a double-blind randomized longitudinal clinical study. I: clinical results. *J Clin Periodontol*. 2009;36(7):604–9. <https://doi.org/10.1111/j.1600-051X.2009.01421.x>. Erratum in: *J Clin Periodontol*. 2009;36(12):1076.
135. Ciancio SG, Lauciello F, Shibly O, Vitello M, Mather M. The effect of an antiseptic mouthrinse on implant maintenance: plaque and peri-implant gingival tissues. *J Periodontol*. 1995;66(11):962–5.

136. Felo A, Shibly O, Ciancio SG, Lauciello FR, Ho A. Effects of subgingival chlorhexidine irrigation on peri-implant maintenance. *Am J Dent*. 1997;10(2):107–10.
137. Groenendijk E, Dominicus JJ, Moorer WR, Aartman IH, van Waas MA. Microbiological and clinical effects of chlorhexidine enclosed in fixtures of 3I-Titamed implants. *Clin Oral Implants Res*. 2004;15(2):174–9.
138. Renvert S, Lessem J, Lindahl C, Svensson M. Treatment of incipient peri-implant Infections using topical minocycline microspheres versus topical chlorhexidine gel as an adjunct to mechanical debridement. *J Int Acad Periodontol*. 2004;6(4 Suppl):154–9.
139. Strooker H, Rohn S, Van Winkelhoff AJ. Clinical and microbiologic effects of chemical versus mechanical cleansing in professional supportive implant therapy. *Int J Oral Maxillofac Implants*. 1998 Nov-Dec;13(6):845–50.
140. Tawse-Smith A, Duncan WJ, Payne AG, Thomson WM, Wennström JL. Relative effectiveness of powered and manual toothbrushes in elderly patients with implant-supported mandibular overdentures. *J Clin Periodontol*. 2002;29(4):275–80.
141. Wolff L, Kim A, Nunn M, Bakdash B, Hinrichs J. Effectiveness of a sonic toothbrush in maintenance of dental implants. A prospective study. *J Clin Periodontol*. 1998;25(10):821–8.
142. Ramberg P, Lindhe J, Botticelli D, Botticelli A. The effect of a triclosan dentifrice on mucositis in subjects with dental implants: a six-month clinical study. *J Clin Dent*. 2009;20(3):103–7.
143. Romeo E, Chiapasco M, Lazza A, Casentini P, Ghisolfi M, Iorio M, Vogel G. Implant-retained mandibular overdentures with ITI implants. *Clin Oral Implants Res*. 2002;13(5):495–501.
144. Enislidis G, Fock N, Millesi-Schobel G, Klug C, Wittwer G, Yerit K, Ewers R. Analysis of Complications following alveolar distraction osteogenesis and implant placement in the partially edentulous mandible. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2005;100(1):25–30.
145. Walton TR. The outcome of implant-supported fixed prostheses from the prosthodontic perspective: proposal for a classification protocol. *Int J Prosthodont*. 1998 Nov-Dec;11(6):595–601.
146. Tarazona B, Vidal-Infer A, Alonso-Arroyo A. Bibliometric analysis of the scientific production in implantology (2009–2013). *Clin Oral Implants Res*. 2017;28(7):864–70.
147. Fogel DB. Factors associated with clinical trials that fail and opportunities for improving the likelihood of success: a review. *Contemp Clin Trials Commun*. 2018;11:156–64.
148. Heneghan C, Goldacre B, Mathani B, Kamal R. Why clinical trial outcomes fail to translate into benefits for patients. *Trials*. 2017;18(1):1–7.
149. French D, Ofec R, Levin L. Long term clinical performance of 10 871 dental implants with up to 22 years of follow-up: a cohort study in 4247 patients. *Clin Implant Dent Relat Res*. 2021;23(3):289–97.
150. Pjetursson BE, Sailer I, Zwahlen M, Hämmerle CH. (2007). A systematic review of the survival and complication rates of all-ceramic and metal-ceramic reconstructions after an observation period of at least 3 years. Part I: Single crowns. *Clin Oral Implants Res* 18(Suppl 3):73–85; *erratum in Clin Oral Implants Res* 19:326–328, 2008.
151. Papaspyridakos P, Chen CJ, Singh M, Weber HP, Gallucci GO. Success criteria in implant dentistry: a systematic review. *J Dent Res*. 2012;91(3):242–8.
152. Gallucci GO, Doughtie CB, Hwang JW, Fiorellini JP, Weber HP. Five-year results of fixed implant-supported rehabilitations with distal cantilevers for the edentulous mandible. *Clin Oral Implants Res*. 2009a;20:601–7.
153. Levi A, Psoter WJ, Agar JR, Reisine ST, Taylor TD. Patient self-reported satisfaction with maxillary anterior dental implant treatment. *Int J Oral Maxillofac Implants*. 2003;18:113–20.
154. Guckes AD, Scurria MS, Shugars DA. A conceptual framework for understanding outcomes of oral implant therapy. *J Prosthet Dent*. 1996;75:633–9.
155. Fuentealba R, Jofré J. Esthetic failure in implant dentistry. *Dent Clin*. 2015;59(1):227–46.
156. Blencowe NS, Strong S, McNair AGK, Brookes ST, Crosby T, Griffin SM, Blazeby JM. Reporting of short-term clinical outcomes after esophagectomy: a systematic review. *Ann Surg*. 2012;255:658–66.
157. Macefield RC, Jacobs M, Korfage IJ, et al. Developing core outcomes sets: methods for identifying and including patient-reported outcomes (PROs). *Trials*. 2014;15:49.
158. Hirsch BR, Califf RM, Cheng SK, Tasneem A, Horton J, et al. Characteristics of oncology clinical trials: insights from a systematic analysis of ClinicalTrials.gov. *JAMA Intern Med*. 2013;173:972–9.
159. Huskisson EC, Wojtulewski JA. Measurement of side effects of Drugs. *Br Med J*. 1974;2(5921):698–9.
160. Myers MG, Cairns JA, Singer J. The consent form as a possible cause of side effects. *Clin Pharmacol Ther*. 1987;42:250–3.
161. The Standards of Reporting Trials Group (SORT). A proposal for structured reporting of randomized controlled trials. *JAMA*. 1994;272:1926–31.
162. Edwards JE, McQuay HJ, Moore RA, Collins SL. Reporting of adverse effects in clinical trials should be improved: lessons from acute postoperative pain. *J Pain Symptom Manag*. 1999;18(6):427–37.
163. Golder S, Loke YK, Wright K, Norman G. Reporting of adverse events in published and unpublished studies of health care interventions: a systematic review. *PLoS Med*. 2016;13(9):e1002127.
164. Eckert SE, Choi YG, Sanchez AR, Koka S. Comparison of dental implant systems: quality of clinical evidence and prediction of 5-year survival. *Int J Oral Maxillofac Implants*. 2005;20:406–15.
165. Williamson PR, Altman DG, Blazeby JM, Clarke M, Devane D, Gargon E, Tugwell P. Developing core outcome sets for clinical trials: issues to consider. *Trials*. 2012;13(1):1–8.
166. Boric K, Dosenovic S, Jelicic Kadic A, Boric M, Jeric M, Puljak L. Efficacy and safety outcomes in systematic reviews of interventions for postoperative pain in children: comparison against the recommended core outcome set. *Pain Med*. 2018;19(11):2316–21.
167. Dosenovic S, Jelicic Kadic A, Jeric M, Boric M, Markovic D, Vucic K, Puljak L. Efficacy and safety outcome domains and outcome measures in systematic reviews of neuropathic pain conditions. *Clin J Pain*. 2018;34(7):674–84.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.