

The role of ambulatory cardiac rehabilitation in improvement of quality of life, anxiety and depression

Rošić, Damir; Krstaić, Goran; Krstaić, Antonija; Brborović, Ognjen; Filipčić, Igor; Mornar Jelavić, Marko

Source / Izvornik: **Psychiatria Danubina, 2020, 32, 496 - 504**

Conference paper / Rad u zborniku

Publication status / Verzija rada: **Published version / Objavljena verzija rada (izdavačev PDF)**

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:127:201508>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International/Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-07-14**



Repository / Repozitorij:

[University of Zagreb School of Dental Medicine Repository](#)



THE ROLE OF AMBULATORY CARDIAC REHABILITATION IN IMPROVEMENT OF QUALITY OF LIFE, ANXIETY AND DEPRESSION

Damir Rosic¹, Goran Krstacic^{2,3,4}, Antonija Krstacic^{3,4,5}, Ognjen Brborovic^{6,7},
Igor Filipic^{3,4,8} & Marko Mornar Jelavic^{2,9}

¹Institute of Emergency Medicine, Rijeka, Croatia

²Institute for Cardiovascular Prevention and Rehabilitation, Zagreb, Croatia

³The Faculty of Medicine, Josip Juraj Strossmayer University of Osijek, Croatia

⁴University of Applied Health Sciences, Zagreb, Croatia

⁵University Hospital Center "Sestre milosrdnice", Clinical Hospital of Traumatology, Zagreb, Croatia

⁶School of Public Health "Andrija Stampar", Zagreb, Croatia

⁷University of Zagreb, School of Medicine, Zagreb, Croatia

⁸Psychiatric Hospital "Sveti Ivan", Zagreb, Croatia

⁹School of Dental Medicine, Zagreb, Croatia

received: 1.4.2020;

revised: 8.7.2020;

accepted: 3.9.2020

SUMMARY

Background: Psychological reactions may adversely affect recovery after major cardiac events. This study investigates the role and frequently negligible importance of ambulatory cardiac rehabilitation (ACR) in improvement of quality of life (QoL), anxiety and depression at patients with various cardiac pathology.

Subjects and methods: This prospective study included subjects treated for acute coronary syndrome (ACS), those with performed elective revascularization, and OTHERS (after valve replacement, implanted pacemaker or other device, with stable heart failure and coronary artery disease). Their anxiety (State Trait Anxiety Inventory (STAI) questionnaire), depression (Beck Depression Inventory (BDI-II) questionnaire) and QoL data (Short Form Health Survey-36 (SF-36) questionnaire, for physical and mental QoL components) were collected initially and after 3-month of ACR.

Results: ACR underwent 170 patients, aged 59 (53-66 years), predominately males (74.7%). At both genders, median duration of ACR was 12 weeks, with reduction of anxiety and depression scores and improvement in almost all components of QoL ($P < 0.05$), except in mental health and bodily pain in males and females, respectively.

After ACS (63.5%), ACR lasted 12 weeks, with reduction of anxiety and depression scores and improvement in all components of QoL ($P < 0.05$). After elective revascularization (14.1%), ACR lasted 12 weeks, with reduction of anxiety score and improvement in almost all components of QoL ($P < 0.05$), except mental health. At OTHERS (22.4%), ACR lasted 4 weeks, with improvement in almost all components of QoL ($P < 0.05$), except mental health; ACR duration negatively correlated with anxiety and depressive scores ($P < 0.05$).

Conclusion: ACR during 3-month results with improvement of anxiety, depression and QoL at patients with various cardiac pathology.

Key words: ambulatory cardiac rehabilitation - anxiety – depression -- quality of life

* * * * *

INTRODUCTION

Cardiovascular diseases (CVD) are the leading cause of death in our country and worldwide as well (Pagidipati et al. 2013, Ćorić et al. 2014). Despite improved treatment of coronary heart disease (CHD), an organized network of primary percutaneous intervention for treatment of acute myocardial infarction (AMI), and consistent application of evidence-based measures of secondary prevention, a significant number of persons still dies from this disease. Data from the Croatian National Institute of Public Health on the number of deaths in 2013 show that AMI is one of the most common single causes of death in Croatia, at 3456 persons (2063

men and 1393 women) or 6.9% of all mortal outcomes, followed by lung cancers (2802 deaths, or 5.6%) and colorectal cancer (2037 deaths or 4%) (Ćorić et al. 2014, Ivanusa et al. 2015). Psychological reactions that cardiac patients may experience (e.g. anxiety and depression) are strong risk factors for CHD and adversely affect recovery after major CHD events (Contractor 2011, Zullo et al. 2017). Moreover, anxiety and depression may increase the risk of future adverse cardiovascular events at patients without apparent cardiovascular pathology (Sanchez-Gonzales et al. 2013, Rieckmann et al. 2013, Slavich & Irwin 2014, Cohen et al. 2015, Vaccarino et al. 2018).

Cardiac rehabilitation (and secondary prevention) services are defined as comprehensive, long term mea-

asures involving medical evaluation, prescribed exercise, cardiac risk factor modification, education, and counseling (Balady et al. 2007, Arena et al. 2012, Mampuya 2012, Dalal et al. 2015, Doimo et al. 2019). These measures are designed to limit the physiological and psychological effects of cardiac illness, reduce the risk for sudden death or re-infarction, control cardiac symptoms, stabilize or reverse the atherosclerotic process, and enhance the psychosocial and vocational status of selected patients counselling (Balady et al. 2007, Arena et al. 2012, Mampuya 2012, Dalal et al. 2015, Doimo et al. 2019).

Cardiac rehabilitation is classified into three phases (Mampuya 2012, Ivanusa et al. 2015). Phase I (inpatient, hospital) is organized in a coronary or cardiac surgery intensive care unit. The shorter hospital stay with modern cardiology treatment makes it difficult to conduct formal inpatient education and training programs. Thus, inpatient cardiac rehabilitation programs are mostly limited to early mobilization to make self-care possible by discharge, and brief counseling about the nature of the illness, the treatment, risk factors management and follow up planning (Mampuya 2012, Ivanusa et al. 2015, Vaccarino et al. 2018).

Phase II (outpatient, ambulatory cardiac rehabilitation (ACR)) takes place in specialized institutions. It is effective in improving of physical activity level, exercise tolerance and quality of life (QoL). Physical activity is gradually increased according to generally accepted principles. This phase is characterized by teamwork (an interdisciplinary and transdisciplinary approach), continuous monitoring by a cardiologist, and telemetric electrocardiogram monitoring during physical activity. It is usually performed in outpatient centers 3-5 times a week over three months. After non-invasive cardiovascular and psychological diagnostics, patients and their families are informed about the state of their disease. They are educated and counseled on the disease as well as further treatment and behavior aimed at encouraging changes in health behavior and reducing negative psychosocial consequences of the disease (Mampuya 2012, Ivanusa et al. 2015, Vaccarino et al. 2018).

Phase III is provided in cardiac patient clubs that are usually organized in rehabilitation centers, and lasts for the rest of patient's life. This phase is characterized by maintenance, and if possible improvement, of existing improvements achieved by cardiovascular rehabilitation, along with intermittent evaluations by internist-cardiologists, exercise stress test, and risk factor assessment (Mampuya 2012, Ivanusa et al. 2015, Vaccarino et al. 2018).

The aim of this prospective study was to investigate the role and frequently negligible importance of ACR in improvement of QoL, anxiety and depression at patients with various cardiac pathology. This is the first such study in the Republic of Croatia.

SUBJECTS AND METHODS

Subjects

This prospective study was performed in Institute for Cardiovascular Prevention and Rehabilitation, in Zagreb, Republic of Croatia. It was approved by the appropriate Institute Ethics Committee and included subjects treated for acute coronary syndrome (ACS), those with performed elective revascularization (percutaneous coronary intervention (PCI), coronary arteries bypass graft (CABG)), and OTHERS (after valve replacement, implanted pacemaker or other device, with stable heart failure and coronary artery disease).

This second phase of cardiac rehabilitation included very low and moderate risk patients with at least 36 sessions and it was performed in accordance with guidelines (Balady et al. 2007). Those with high risk (unstable angina, decompensated heart failure, complex ventricular arrhythmias, pulmonary pressure ≥ 60 mmHg, intracavitary thrombus, recent thrombophlebitis with or without pulmonary embolism, severe obstructive cardiomyopathies, severe or symptomatic aortic stenosis, uncontrolled inflammatory or infectious pathologies and any musculoskeletal condition that prohibits physical exercise) were excluded. Most patients were referred by their cardiologist, internist, family physician, or cardiac surgeon.

Methods

Instruments

Between February and September 2013, we collected baseline demographic, CHD risk factors and psychosocial functioning data. Demographic data included age, gender and educational level. CHD risk factors data included sedentary lifestyle, obesity, arterial hypertension, dyslipidemia, hyperglycemia/diabetes and smoking. Psychosocial functioning was evaluated with State Trait Anxiety Inventory (STAI) for anxiety, Beck Depression Inventory (BDI-II) for depression, and Short Form Health Survey-36 (SF-36) questionnaire for QoL.

A number of reliability and validity tests have been conducted on the STAI and have provided sufficient evidence that the STAI is an appropriate for anxiety studies in research and clinical settings (Spielberger et al. 1970, Spielberger et al. 1983). It is composed of 20 items. Respondents were encouraged to report their general feelings about anxiety on a four-point scale: 1 (almost never), 2 (sometimes), 3 (often), and 4 (almost always). The total score reveals the anxiety level, ranging from 20 to 80. High scores indicate a high level of trait anxiety; scores of 20-29 indicated mild anxiety, and scores ≥ 30 indicated moderate/severe anxiety.

BDI questionnaire is a 21-item measure of depressive symptoms (Beck et al. 1996, Ceccarini et al. 2014). For each item, participants chose the statement that best reflected how they felt. Responses to the individual

items were then summed, with scores ranging from 0 to 63. Scores of 0 to 9 indicated minimal depression, scores of 10 to 18 indicated mild depression, and scores of 19 to 63 indicated moderate to severe depressive symptoms. It is a relevant psychometric instrument, showing high reliability, capacity to discriminate between depressed and non-depressed subjects, and improved concurrent, content, and structural validity. Based on available psychometric evidence, it can be viewed as a cost-effective questionnaire for measuring the severity of depression, with broad applicability for research and clinical practice worldwide (Wang & Gorenstein 2013).

The SF-36 is a generic measure of health status comprising 36 items related to eight dimensions: physical functioning for the limitation in performing all physical activities, role physical for problems with work or other daily activities, bodily pain, general health, vitality, social functioning, role emotional and mental health (McHorney et al. 1994). The dimensions such as physical functioning, role physical, bodily pain and general health reflect the physical component of health, while the dimensions vitality, social functioning, role emotional and mental health reflect the mental components of health. SF-36 has been adapted and translated into several languages, and its validity and reliability established in several countries (Bullinger et al. 1998, Gandek et al. 1998, Wagner et al. 1998).

ACR program

After an initial analysis, we created an individualized rehabilitation plan (the type, intensity, duration, and frequency of cardiorespiratory therapy), and the patient was placed into one of five existing intensity level groups for the treatment during 12 weeks. Education of the patient and family/partner was performed by all members of the rehabilitation team both individually and in groups, using advice, workshops, and lectures from the fields of cardiology, physical medicine and rehabilitation, psychology, and psychiatry. The psychologist performed patient counseling, education, group therapy based on behavioral-cognitive principles, and participating in cardiovascular consultation. The contents and scope of the counseling, conducted during the interview, and if needed in additional weekly sessions, depended on the patients, their primary disease, comorbidities, and possible psychological difficulties. The psychologist gave two lectures on psychological risk factors for CHD as part of the regular lecture cycle for patients and their families.

We offered to patients to take part in a small, closed relaxation and weight-reduction groups. They learned about diaphragmatic breathing, facilitating relaxation, adequate communication skills, anger management, assertiveness, problem-solving schemes, recognizing and coping with manipulation, etc. The weight reduction group focused on the control of the stimuli which lead to eating, and the control of alimentary processes. Gra-

duel reduction of daily energy intake values coupled with proper ingredient choices and combinations depended on individual risk factors. Both groups met once a week. The psychologist took part in cardiovascular consultation and informed other members on the psychological risk factors and characteristics of the particular patient's personality.

Data collection after ACR

After 3-month of ACR, we performed control measurements of the all above mentioned baseline parameters.

Statistical analysis

Qualitative data are presented with absolute number and percentage. We used McNemar's test of paired proportions for its analysis. Quantitative data with normal distribution are presented with mean and standard deviation (SD), and compared with paired samples t-test; data without normal distribution are presented with median and corresponding interquartile range.

Correlations between the variables were investigated with Pearson's r , Spearman's ρ and Kendall's τ correlation coefficients, classified as a very weak (0-0.19), weak (0.20-0.39), moderate (0.40-0.59), strong (0.60-0.79) and very strong (0.80-1).

The level of statistical significance was set at $P < 0.05$. Processing was done with MedCalc 12.7.0.0 for Windows software.

RESULTS

Of the total of 170 patients, aged 59 (53-66) years, we had the highest prevalence of males (74.7%). ACR was the most frequently indicated after ACS (63.5%), then elective revascularization (14.1%) and at *others* (22.4%). We have found the highest prevalence of sedentary lifestyle (88.2%), then obesity (74.1%), hypertension (64.7%), smoking (48.2%), dyslipidemia (40.6%) and hyperglycemia/diabetes (34.7%). The most of patients had middle (65.3%), then high (24.1%) and low (10.6%) level of education. We got the following results:

At overall patients

ACR lasted 12 (8-12) weeks, with reduction of mean values of anxiety (39.2 vs 37.1) and depression scores (10.4 vs 9.9) ($P < 0.05$), without significant differences in the prevalence of mild (43 (25.3%) vs 43 (25.3%), $P = 1.000$) and moderate/severe anxiety (127 (74.7%) vs 127 (74.7%), $P = 1.000$), and minimal (95 (55.9%) vs 97 (57.1%), $P = 0.500$), mild (46 (27.1%) vs 48 (28.2%), $P = 0.688$) and moderate/severe depression (29 (17.1%) vs 25 (14.7%), $P = 0.125$) (Figure 1, Table 1). QoL has improved in mean values of general health (62.0 vs 72.0), physical (61.3 vs 70.1) and emotional role (69.4 vs 77.8), physical (75.9 vs 80.1) and social functioning (76.3 vs 79.3), bodily pain (76.6 vs 79.0), vitality (59.3 vs 61.6) and mental health (65.1 vs 66.4) ($P < 0.05$) (Table 1).




Figure 1. The number of patients with various level of anxiety and depression before (A) and after (B) ACR (N=170), without significant differences (McNemar's test, statistical significance with $P < 0.05$; ACR – ambulatory cardiac rehabilitation)

Table 1. Differences between the examined data before and after ACR at overall patients (N=170)

Findings	Parameters	Before ACR (mean)	SD	After ACR (mean)	SD	t	df	P
Psychological testing	Anxiety score	39.2	12.2	37.1	9.8	-6.123	169	0.000
	Depression score	10.4	8.6	9.9	7.6	-3.888	169	0.000
QoL (Physical components)	Physical functioning	75.9	21.4	80.1	15.7	7.713	169	0.000
	Role physical	61.3	39.3	70.1	30.8	9.601	169	0.000
	Bodily pain	76.6	25.5	79.0	22.3	4.703	169	0.000
	General health	62.0	20.3	72.0	12.5	11.448	169	0.000
QoL (Mental components)	Vitality (energy/fatigue)	59.3	16.7	61.6	14.2	7.183	169	0.000
	Social functioning	76.3	23.0	79.3	18.9	5.066	169	0.000
	Role emotional	69.4	39.1	77.8	30.1	6.180	169	0.000
	Mental health	65.1	15.9	66.4	12.5	2.588	169	0.011

ACR – ambulatory cardiac rehabilitation; QoL – quality of life; Data presented with mean and standard deviation (SD) were compared with paired t-test. Statistical significance with $P < 0.05$

Values of anxiety and depression scores negatively correlated with values of QoL parameters before (anxiety: $r = -0.27$ to -0.64 ; depression: $r = -0.21$ to -0.61) and after ACR (anxiety: $r = -0.23$ to -0.56 ; depression: $r = -0.20$ to -0.50) ($P < 0.05$). It was weaker after ACR, but still present. Depression positively correlated with anxiety scores at both time points ($r_1 = 0.65$, $r_2 = 0.57$) ($P < 0.05$).

At males

ACR lasted 12 (8-12) weeks, with reduction of mean values of anxiety (39.1 vs 37.0) and depression scores (9.8 vs 9.2) ($P < 0.05$), without significant differences in the prevalence of mild (34 (26.8%) vs 35 (27.6%), $P = 1.000$) and moderate/severe anxiety (93 (73.2%) vs 92 (72.4%), $P = 1.000$), and minimal (80 (63.0%) vs 82 (64.6%), $P = 0.500$), mild (26 (20.5%) vs 26 (20.5%), $P = 1.000$) and moderate/severe depression (21 (16.5%) vs 19 (15.0%), $P = 0.500$) (Table 2). QoL has improved in mean values of general health (62.2 vs 72.5), physical (64.8 vs 72.6) and emotional role (68.8 vs 77.4),

physical (77.7 vs 81.7) and social functioning (77.5 vs 80.1), bodily pain (77.4 vs 80.0) and vitality (61.2 vs 63.5) ($P < 0.05$), without improvement of mental health (66.8 vs 67.9) (Table 2).

At females

ACR lasted 12 (6-12) weeks, with reduction of mean values of anxiety (39.6 vs 37.3) and depression scores (12.1 vs 11.7) ($P < 0.05$), without significant differences in the prevalence of mild (9 (20.9%) vs 8 (18.6%), $P = 1.000$) and moderate/severe anxiety (34 (79.1%) vs 35 (81.4%), $P = 1.000$), and minimal (15 (34.9%) vs 15 (34.9%), $P = 1.000$), mild (20 (46.5%) vs 22 (51.2%), $P = 0.500$) and moderate/severe depression (8 (18.6%) vs 6 (14.0%), $P = 0.500$) (Table 3). QoL has improved in mean values of general health (61.3 vs 70.6), physical (51.2 vs 62.8) and emotional role (71.3 vs 79.1), physical (70.3 vs 75.3) and social functioning (72.7 vs 76.7), vitality (53.5 vs 56.3) and mental health (60.3 vs 62.0) ($P < 0.05$), without improvement of bodily pain (74.4 vs 76.3) (Table 3).

Table 2. Differences between the examined data before and after ACR at males (N=127)

Findings	Parameters	Before ACR (mean)	SD	After ACR (mean)	SD	t	df	P
Psychological testing	Anxiety score	39.1	12.5	37.0	10.2	-5.252	126	0.000
	Depression score	9.8	9.0	9.2	7.8	-3.301	126	0.001
QoL (Physical components)	Physical functioning	77.7	21.3	81.7	15.6	6.300	126	0.000
	Role physical	64.8	38.7	72.6	30.4	7.611	126	0.000
	Bodily pain	77.4	25.1	80.0	21.5	4.275	126	0.000
	General health	62.2	21.5	72.5	13.0	9.948	126	0.000
QoL (Mental components)	Vitality (energy/fatigue)	61.2	17.2	63.5	14.5	5.530	126	0.000
	Social functioning	77.5	22.3	80.1	18.7	3.998	126	0.000
	Role emotional	68.8	39.6	77.4	30.5	5.446	126	0.000
	Mental health	66.8	15.7	67.9	11.7	1.824	126	0.071

ACR – ambulatory cardiac rehabilitation; QoL – quality of life; Data presented with mean and standard deviation (SD) were compared with paired t-test. Statistical significance with P<0.05

Table 3. Differences between the examined data before and after ACR at females (N=43)

Findings	Parameters	Before ACR (mean)	SD	After ACR (mean)	SD	t	df	P
Psychological testing	Anxiety score	39.6	11.5	37.3	8.8	-3.112	42	0.003
	Depression score	12.1	7.3	11.7	6.7	-2.324	42	0.025
QoL (Physical components)	Physical functioning	70.3	21.1	75.3	15.3	4.480	42	0.000
	Role physical	51.2	39.7	62.8	31.0	6.043	42	0.000
	Bodily pain	74.4	26.8	76.3	24.7	1.965	42	0.056
	General health	61.3	16.3	70.6	11.2	5.623	42	0.000
QoL (Mental components)	Vitality (energy/fatigue)	53.5	13.9	56.3	12.0	5.225	42	0.000
	Social functioning	72.7	24.7	76.7	19.6	3.138	42	0.003
	Role emotional	71.3	38.2	79.1	29.1	2.892	42	0.006
	Mental health	60.3	15.5	62.0	13.8	2.301	42	0.026

ACR – ambulatory cardiac rehabilitation; QoL – quality of life; Data presented with mean and standard deviation (SD) were compared with paired t-test. Statistical significance with P<0.05

At patients after ACS

ACR lasted 12 (11.5-12) weeks, with reduction of mean values of anxiety (39.9 vs 37.3) and depression scores (10.7 vs 10.1) (P<0.05), without significant differences in the prevalence of mild (24 (22.2%) vs 27 (25.0%), P=0.375) and moderate/severe anxiety (84 (77.8%) vs 81 (75.0%), P=0.375), and minimal (61 (56.5%) vs 62 (57.4%), P=1.000), mild (28 (25.9%) vs 31 (28.7%), P=0.625) and moderate/severe depression (19 (17.6%) vs 15 (13.9%), P=0.125) (Table 4). QoL has improved in mean values of general health (61.3 vs 71.3), physical (64.1 vs 71.3) and emotional role (66.0 vs 75.0), physical (78.4 vs 81.9) and social functioning (75.6 vs 79.3), bodily pain (78.7 vs 80.9), vitality (59.1 vs 61.8) and mental health (63.5 vs 65.4) (P<0.05) (Table 4).

At patients after elective revascularization

ACR lasted 12 (11.5-12) weeks, with reduction of mean values of anxiety score (39.8 vs 37.8), without significant differences in mean values of depression score (10.5 vs 10.5) and prevalence of mild (8 (33.3%) vs 6 (25.0%), P=0.250) and moderate/severe anxiety (16 (66.7%) vs 18 (75.0%), P=0.250), and minimal (11 (45.8%) vs 11 (45.8%), P=1.000), mild (9 (37.5%) vs 9 (37.5%), P=1.000) and moderate/severe depression (4

(16.7%) vs 4 (16.7%), P=1.000) (Table 5). QoL has improved in mean values of general health (61.0 vs 71.2), physical (44.8 vs 60.4) and emotional role (62.5 vs 70.8), physical (64.6 vs 72.7) and social functioning (70.8 vs 72.9), bodily pain (67.5 vs 71.2) and vitality (59.6 vs 61.5), without improvement of mental health (66.5 vs 67.0) (P<0.05) (Table 5).

At others

ACR lasted 4 (4-12) weeks, without significant differences in mean values of anxiety (36.9 vs 35.9) and depression scores (9.3 vs 8.7), and prevalence of mild (11 (28.9%) vs 10 (26.3%), P=1.000) and moderate/severe anxiety (27 (71.1%) vs 28 (73.7%), P=1.000), and minimal (23 (60.5%) vs 24 (63.2%), P=1.000), mild (9 (23.7%) vs 8 (21.1%), P=1.000) and moderate/severe depression (6 (15.8%) vs 6 (15.8%), P=1.000) (Table 6). QoL has improved in mean values of general health (64.6 vs 74.8), physical (63.8 vs 73.0) and emotional role (83.3 vs 90.4), physical (75.8 vs 79.6) and social functioning (81.6 vs 83.2), bodily pain (76.6 vs 78.3) and vitality (59.5 vs 61.4) (P<0.05), but without improvement of mental health (68.8 vs 68.8) (Table 6). In this group, we had negative correlation of ACR duration with anxiety ($\rho=-0.31$, P=0.058; $\tau=-0.25$, P=0.024) and depressive scores ($\rho=-0.29$, P=0.078; $\tau=-0.23$, P=0.039).

Table 4. Differences between the examined data before and after ACR at patients after ACS (N=108)

Findings	Parameters	Before ACR (mean)	SD	After ACR (mean)	SD	t	df	P
Psychological testing	Anxiety score	39.9	12.0	37.3	9.3	-5.538	107	0.000
	Depression score	10.7	8.7	10.1	7.5	-3.586	107	0.001
QoL (Physical components)	Physical functioning	78.4	19.5	81.9	14.7	5.711	107	0.000
	Role physical	64.1	39.3	71.3	31.1	6.563	107	0.001
	Bodily pain	78.7	25.6	80.9	22.3	3.471	107	0.001
	General health	61.3	20.4	71.3	12.7	8.987	107	0.000
QoL (Mental components)	Vitality (energy/fatigue)	59.1	17.1	61.8	14.2	5.705	107	0.000
	Social functioning	75.6	24.2	79.3	19.2	4.237	107	0.000
	Role emotional	66.0	40.2	75.0	31.0	5.167	107	0.000
	Mental health	63.5	15.1	65.4	11.7	2.917	107	0.004

ACR – ambulatory cardiac rehabilitation; QoL – quality of life; Data presented with mean and standard deviation (SD) were compared with paired t-test. Statistical significance with P<0.05

Table 5. Differences between the examined data before and after ACR at patients after elective revascularization (N=24)

Findings	Parameters	Before ACR (mean)	SD	After ACR (mean)	SD	t	df	P
Psychological testing	Anxiety score	39.8	13.4	37.8	11.5	-2.470	23	0.021
	Depression score	10.5	8.4	10.5	8.0	0.171	23	0.866
QoL (Physical components)	Physical functioning	64.6	25.7	72.7	17.6	4.175	23	0.000
	Role physical	44.8	36.1	60.4	27.5	6.191	23	0.000
	Bodily pain	67.5	26.5	71.2	21.7	2.415	23	0.024
	General health	61.0	19.9	71.2	12.0	4.392	23	0.000
QoL (Mental components)	Vitality (energy/fatigue)	59.6	18.1	61.5	15.9	2.584	23	0.017
	Social functioning	70.8	23.2	72.9	20.4	2.145	23	0.043
	Role emotional	62.5	39.7	70.8	34.5	2.304	23	0.031
	Mental health	66.5	19.6	67.0	16.1	0.430	23	0.671

ACR – ambulatory cardiac rehabilitation; QoL – quality of life; Data presented with mean and standard deviation (SD) were compared with paired t-test. Statistical significance with P<0.05

Table 6. Differences between the examined data before and after ACR at OTHERS (N=38)

Findings	Parameters	Before ACR (mean)	SD	After ACR (mean)	SD	t	df	P
Psychological testing	Anxiety score	36.9	12.2	35.9	10.1	-1.586	37	0.121
	Depression score	9.3	8.8	8.7	7.6	-1.981	37	0.055
QoL (Physical components)	Physical functioning	75.8	22.1	79.6	16.3	3.399	37	0.002
	Role physical	63.8	39.7	73.0	31.5	4.646	37	0.000
	Bodily pain	76.6	23.8	78.3	22.3	2.197	37	0.034
	General health	64.6	20.3	74.8	12.4	5.435	37	0.000
QoL (Mental components)	Vitality (energy/fatigue)	59.5	14.9	61.4	13.5	3.812	37	0.001
	Social functioning	81.6	18.3	83.2	16.3	2.368	37	0.023
	Role emotional	83.3	32.7	90.4	20.4	2.458	37	0.019
	Mental health	68.8	15.3	68.8	12.0	0.000	37	1.000

ACR – ambulatory cardiac rehabilitation; QoL – quality of life; Data presented with mean and standard deviation (SD) were compared with paired t-test. Statistical significance with P<0.05

DISCUSSION

According to the literature data, about 20% patients with CHD or heart failure is depressed, which is ≥ 3 times greater than in the general population (Lane et al. 2002, Ruo et al. 2003, Rutledge et al. 2006, Cohen et al. 2015, Vaccarino et al. 2018). Patients with CHD, heart failure and depressive symptoms are more likely to have physical limitations and poor QoL (Ruo et al. 2003, Gottlieb et al. 2009, Cohen et al. 2015, Vaccarino et al.

2018). Moreover, they have increased risk for recurrent cardiovascular events and mortality; depressed patients with ACS have double the risk of future adverse cardiovascular events, which is similar to conventional risk factors such as reduced ejection fraction and diabetes (Jiang et al. 2001, Nicholson et al. 2006, Bartoli et al. 2013, Cohen et al. 2015, Vaccarino et al. 2018). Depression may be associated with lower physical activity, more intensive smoking and lower adherence to cardiovascular medications (Kronish et al. 2006, Bautista et al. 2012, Cohen et

al. 2015, Vaccarino et al. 2018). It has been linked with inflammatory processes, autonomic nervous system dysfunction, and impaired coronary flow reserve that increases risk of myocardial ischemia (Sanchez-Gonzales et al. 2013, Rieckmann et al. 2013, Slavich & Irwin 2014, Cohen et al. 2015, Vaccarino et al. 2018).

Anxiety is associated with cigarette smoking, excess alcohol consumption, lower physical activity, and poor diet, which increases the risk of CVD (Strine et al. 2005, Antonogeorgos et al. 2012, Cohen et al. 2015, Celano et al. 2016). Anxiety leads to excess activation of the hypothalamic-pituitary-adrenal axis and sympathetic nervous system, increases release of plasma catecholamines, and causes endothelial damage, ultimately leading to atherosclerosis, CHD, and acute coronary events (Cohen et al. 2015, Celano et al. 2016). Acute anxiety is associated with increased cardiovascular reactivity to stress and resting heart rate, decreased heart rate variability, baroreflex dysfunction, and greater variability in ventricular repolarization (Rozanski et al. 2005, Cohen et al. 2015, Celano et al. 2016). These effects may increase the risk of incident CVD and lower the threshold for cardiac ischemia, arrhythmias, and sudden cardiac death.

Generally, cardiovascular rehabilitation has been proven to be an effective treatment to continuously apply after pharmacological and invasive and/or surgical treatment of the acute phase of the disease (e.g. after AMI) (Balady et al. 2007, Mampuya 2012, Arena et al. 2012, Dalal et al. 2015, Doimo et al. 2019). ACR allows clinically stable (low to moderate risk) patients to lead active lives in their places of residence, thanks to the efforts of a rehabilitation team of medical workers (Arena et al. 2012). The psychologists have special role in rehabilitation process through psychodiagnostics, patient counseling, education, group therapy based on behavioral-cognitive principles, and participating in cardiovascular consultation. Psychodiagnostics includes two types of activity – testing and conversation/interviews. In Croatia, as in other countries, it is adaptable and economical, but underused because it is ignored by medical professionals themselves, but also due to possible remoteness of available rehabilitation centers. This participation is particularly poor in rural areas and in eligible patients who have lower socioeconomic status, limited education, advanced age, and/or female sex (Arena et al. 2012, Ivanusa et al. 2015).

According to our knowledge, this is the first study in the Republic of Croatia and literature about the effectiveness of ACR on improvement of psychological, QoL and cardiac status at patients with various spectrum of cardiac pathology. Also, the literature data about the psychosocial effectiveness of ACR after ACS, revascularization and, especially, after valve replacement and device implantation procedures is somewhat scarce or insufficient. These facts represent the unique characteristics and advantages of our study.

We have found improvement of anxiety and depression scores, as well as of QoL, mostly after ACS and elective revascularization, which could be explained with longer duration of ACR in these groups. Also, correlation between anxiety, depression and QoL was weaker after rehabilitation, but it was still present. As no one of the patients have taken no medications for anxiety and depression before and during the rehabilitation procedure, that finding may suggest successful rehabilitation. So, we are sure that the additional efforts of psychologists, as well as an additional involvement of psychiatrists, may lead to much more improvement of anxiety, depression and QoL.

Cardiac rehabilitation in distressed patients may be difficult (McGrady et al. 2009, Contractor 2011, Zullo 2017), but reduces the blood pressure, blood glucose, lipid levels and body weight (central obesity and metabolic syndrome), which are traditional risk factors for myocardial (re)infarction due to atherosclerosis, chronic inflammation and endothelial dysfunction (Yusuf et al. 2004, Ranjit et al. 2007, Mornar Jelavic et al. 2015).

After an acute coronary event, elective percutaneous coronary revascularization, or CABG, conduction of ACR during two and 3-month results with improvement of anxiety and depressive scores, and QoL (Milani et al. 1996, Tuniz et al. 2004, Pourafkari et al. 2016). Other authors also reported effectiveness of two and 3-month cardiac rehabilitation at patients with CHD and after CABG in improvement of QoL and depression (Solak et al. 2015).

At patients with stable heart failure (NYHA class II-III), ACR during 2-month may result with better values of anxiety, depression and QoL (Kulcu et al. 2007, Keihani et al. 2015).

Study limitations

The sample of patients who participated in the study was rather specific, due to the fact that we excluded high risk subjects at whom ACR was contraindicated. Furthermore, there was no parallel control group of patients who did not participate in the rehabilitation program, so the resulting effects cannot be clearly attributed to the effectiveness of the ACR. Also, there was no follow-up that would test the continuance of the observed changes. So, future investigations should be performed on a larger number of patients with monitoring of their psychosocial and cardiovascular condition during follow up period, for at least 6-12 months.

CONCLUSION

The findings of this study showed that ACR is associated with improvement of anxiety, depression and QoL at patients with various cardiac pathology. We emphasized the necessity of more intensive psychological approach, as well as possible future cooperation

with psychiatrists, given the known negative effects of depression and anxiety on the QoL and recovery of patients after major CHD events. Finally, we do hope that this article can serve as an incentive for future scientific research and more frequent application of ACR as a safe and successful procedure for patients with various cardiac pathology.

Acknowledgements: None.

Conflict of interest: None to declare.

Contribution of individual authors:

Damir Rosic, Goran Krstacic, Igor Filipic & Marko Mornar Jelavic: design study, analyses of data results, statistical analyses, literature searches, manuscript writing.

Antonija Krstacic & Ognjen Brborovic: design study, analyses of data results, statistical analyses, literature searches.:

References

1. Antonogeorgos G, Panagiotakos DB, Pitsavos C, Papageorgiou C, Chrysohoou C, Papadimitriou GN, et al.: Understanding the role of depression and anxiety on cardiovascular disease risk, using structural equation modeling; the mediating effect of the Mediterranean diet and physical activity: the ATTICA study. *Ann Epidemiol* 2012; 22:630-7
2. Arena R, Williams M, Forman DE, Cahalin LP, Coke L, Myers J, et al.: American Heart Association Exercise, Cardiac Rehabilitation and Prevention Committee of the Council on Clinical Cardiology, Council on Epidemiology and Prevention, and Council on Nutrition, Physical Activity and Metabolism. Increasing referral and participation rates to outpatient cardiac rehabilitation: the valuable role of healthcare professionals in the inpatient and home health settings: a science advisory from the American Heart Association. *Circulation* 2012; 125:1321-9
3. Balady GJ, Williams MA, Ades PA, Bittner V, Comoss P, Foody JM, et al.: Core components of cardiac rehabilitation/ secondary prevention programs: 2007 Update. A scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation* 2007; 115:2675-82
4. Bartoli F, Lillia N, Lax A, Crocamo C, Mantero V, Carrà G, et al.: Depression after stroke and risk of mortality: a systematic review and meta-analysis. *Stroke Res Treat* 2013; 2013:862978
5. Bautista LE, Vera-Cala LM, Colombo C, Smith P: Symptoms of depression and anxiety and adherence to antihypertensive medication. *Am J Hypertens* 2012; 25:505-11
6. Beck AT, Steer RA, Ball R, Ranieri W: Comparison of Beck Depression Inventories -IA and -II in psychiatric outpatients. *J Pers Assess* 1996; 67:588-97
7. Bullinger M, Alonso J, Apolone G, Lepège A, Sullivan M, Wood-Dauphinee S, et al.: Translating health status questionnaires and evaluating their quality: The IQOLA project approach. *J Clin Epidemiol* 1998; 51:913-23
8. Ceccarini M, Manzoni GM, Castelnovo G: Assessing depression in cardiac patients: what measures should be considered? *Depress Res Treat* 2014; 2014:148256
9. Celano CM, Daunis DJ, Lokko HN, Campbell KA, Huffman JC: Anxiety Disorders and Cardiovascular Disease. *Curr Psychiatry Rep* 2016; 18:101
10. Cohen BE, Edmondson D, Kronish IM: State of the Art Review: Depression, Stress, Anxiety, and Cardiovascular Disease. *Am J Hypertens* 2015; 28:1295-302
11. Contractor AS: Cardiac rehabilitation after myocardial infarction. *J Assoc Physicians India* 2011; 59:51-5
12. Čorić T, Miler A: Izvješće o umrlim osobama u Hrvatskoj u 2013. godini. Hrvatski zavod za javno zdravstvo, 2014
13. Dalal HM, Doherty P, Taylor RS: Cardiac rehabilitation. *BMJ* 2015; 351:h5000
14. Doimo S, Fabris E, Piepoli M, Barbati G, Antonini-Canterin F, Bernardi G, et al.: Impact of ambulatory cardiac rehabilitation on cardiovascular outcomes: a long-term follow-up study. *Eur Heart J* 2019; 40:678-685
15. Gandek B, Ware JE Jr, Aaronson NK, Alonso J, Apolone G, Bjorner J, et al.: Tests of data quality, scaling assumptions, and reliability of the SF-36 in eleven countries: Results from the IQOLA project. *International Quality of Life Assessment*. *J Clin Epidemiol* 1998; 51:1149-58
16. Gottlieb SS, Kop WJ, Ellis SJ, Binkley P, Howlett J, O'Connor C, et al.: HF-ACTION Investigators. Relation of depression to severity of illness in heart failure (from Heart Failure And a Controlled Trial Investigating Outcomes of Exercise Training [HF-ACTION]). *Am J Cardiol* 2009; 103:1285-9
17. Ivanusa M, Narancic SK, Glavas VD, Kruhek LD, Heinrich M, Mazuran BL, et al.: Outpatient Cardiovascular Rehabilitation in Croatia. *Cardiol Croat* 2015; 10:28-42
18. Jiang W, Alexander J, Christopher E, Kuchibhatla M, Gaulden LH, Cuffe MS, et al.: Relationship of depression to increased risk of mortality and rehospitalization in patients with congestive heart failure. *Arch Intern Med* 2001; 161:1849-56
19. Keihani D, Kargarfard D, Mokhtari M: Cardiac effects of exercise rehabilitation on quality of life, depression and anxiety in patients with heart failure. *Journal of Fundamentals of Mental Health* 2015; 17:13-9
20. Kronish IM, Rieckmann N, Halm EA, Shimbo D, Vorchheimer D, Haas DC, et al.: Persistent depression affects adherence to secondary prevention behaviors after acute coronary syndromes. *J Gen Intern Med* 2006; 21:1178-83
21. Kulcu DG, Kurtai Y, Tur BS, Gülec S, Seekin B: The effect of cardiac rehabilitation on quality of life, anxiety and depression in patients with congestive heart failure. A randomized controlled trial, short-term results. *Eura Medicophys* 2007; 43:489-97
22. Lane D, Carroll D, Ring C, Beevers DG, Lip GY: The prevalence and persistence of depression and anxiety following myocardial infarction. *Br J Health Psychol* 2002; 7:11-21

23. Mampuya WM: Cardiac rehabilitation past, present and future: an overview. *Cardiovasc Diagn Ther* 2012; 2:38-49
24. McHorney CA, Ware JE Jr, Lu JF, Sherbourne CD: The MOS 36-item Short-Form Health Survey (SF-36): III. Tests of data quality, scaling assumptions, and reliability across diverse patient groups. *Med Care* 1994; 32:40-66
25. McGrady A, McGinnis R, Badenhop D, Bentle M, Rajput M: Effects of depression and anxiety on adherence to cardiac rehabilitation. *J Cardiopulm Rehabil Prev* 2009; 29:358-64
26. Milani RV, Lavie CJ, Cassidy MM: Effects of cardiac rehabilitation and exercise training programs on depression in patients after major coronary events. *Am Heart J* 1996; 132:726-32
27. Mornar Jelavic M, Babic Z, Pintaric H: Metabolic syndrome: influence on clinical severity and prognosis in patients with acute ST-elevation myocardial infarction treated with primary percutaneous coronary intervention. *Acta Cardiol* 2015; 70:149-56
28. Nicholson A, Kuper H, Hemingway H: Depression as an aetiological and prognostic factor in coronary heart disease: a meta-analysis of 6362 events among 146 538 participants in 54 observational studies. *Eur Heart J* 2006; 27:2763-74
29. Pagidipati NJ, Gaziano TA: Estimating deaths from cardiovascular disease: a review of global methodologies of mortality measurement. *Circulation* 2013; 127:749-56
30. Pourafkari L, Ghaffari S, Tajlil A, Shahamfar J, Hedayati S, Nader DA: The impact of cardiac rehabilitation program on anxiety and depression levels after coronary artery bypass graft surgery. *Cor et Vasa* 2016; e384-e390
31. Ranjit N, Diez-Roux AV, Shea S, Cushman M, Seeman T, Jackson SA et al.: Psychosocial factors and inflammation in the multi-ethnic study of atherosclerosis. *Arch Intern Med* 2007; 167:174-81
32. Rieckmann N, Kronish IM, Shapiro PA, Whang W, Davidson KW: Serotonin reuptake inhibitor use, depression, and long-term outcomes after an acute coronary syndrome: a prospective cohort study. *JAMA Intern Med* 2013; 173:11501
33. Rozanski A, Blumenthal JA, Davidson KW, Saab PG, Kubzansky L: The epidemiology, pathophysiology, and management of psychosocial risk factors in cardiac practice: the emerging field of behavioral cardiology. *J Am Coll Cardiol* 2005; 45:637-51
34. Ruo B, Rumsfeld JS, Hlatky MA, Liu H, Browner WS, Whooley MA: Depressive symptoms and health-related quality of life: the Heart and Soul Study. *JAMA* 2003; 290:215-21
35. Rutledge T, Reis VA, Linke SE, Greenberg BH, Mills PJ: Depression in heart failure a meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes. *J Am Coll Cardiol* 2006; 48:1527-37
36. Sanchez-Gonzalez MA, May RW, Koutnik AP, Kabbaj M, Fincham FD: Sympathetic vasomotor tone is associated with depressive symptoms in young females: a potential link between depression and cardiovascular disease. *Am J Hypertens* 2013; 26:1389-97
37. Slavich GM, Irwin MR: From stress to inflammation and major depressive disorder: a social signal transduction theory of depression. *Psychol Bull* 2014; 140:774-815
38. Solak Ö, Yaman F, Ulaşlı AM, Eroğlu S, Akçi Ö, Özkeçeci G, et al.: Improvement in Life Quality, Functional Capacity and Depression Level After Cardiac Rehabilitation. *Turk J Phys Med Rehab* 2015; 61:130-5
39. Spielberger CD, Gorsuch RL, Lushene RE: *STAI Manual for the State-Trait Anxiety Inventory*. Consulting Psychologist Press, Palo Alto, CA, 1970
40. Spielberger CD: *State-Trait Anxiety Inventory for Adults. Sampler Set, Manual Set, Scoring Key*. Consulting Psychologists Press, Palo Alto CA, 1983
41. Strine TW, Chapman DP, Kobau R, Balluz L: Associations of self-reported anxiety symptoms with health-related quality of life and health behaviors. *Soc Psychiatry Psychiatr Epidemiol* 2005; 40:432-8
42. Tuniz D, Bernardi G, Molinis G, Valente M, D'Odorico N, Mirolo R, et al.: Ambulatory cardiac rehabilitation with individualized care after elective coronary angioplasty. *Eur Heart J Suppl* 2004; 6:J37-J46
43. Vaccarino V, Badimon L, Bremner JD, Cenko E, Cubedo J, Dorobantu M, et al.: ESC Scientific Document Group Reviewers. Depression and coronary heart disease: 2018 ESC position paper of the working group of coronary pathophysiology and microcirculation developed under the auspices of the ESC Committee for Practice Guidelines. *Eur Heart J* 2019
44. Wagner AK, Gandek B, Aaronson NK, Acquadro C, Alonso J, Apolone G, et al.: Cross-cultural comparisons of the content of SF-36 translations across 10 countries: Results from the IQOLA project. *International Quality of Life Assessment*. *J Clin Epidemiol* 1998; 51:925-32
45. Wang YP, Gorenstein C: Psychometric properties of the Beck Depression Inventory-II: a comprehensive review. *Rev Bras Psiquiatr* 2013; 35:416-31
46. Yusuf S, Hawken S, Ounpuu S, Dans T, Avezum A, Lanas F, et al.: Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004; 364:937-52
47. Zullo MD, Gathright EC, Dolansky MA, Josephson RA, Cheruvu VK, Hughes JW: Influence of Depression on Utilization of Cardiac Rehabilitation Postmyocardial Infarction: A study of 158 991 medicare beneficiaries. *J Cardiopulm Rehabil Prev* 2017; 37:22-29

Correspondence:

Marko Mornar Jelavic, MD, PhD
Institute for Cardiovascular Prevention and Rehabilitation
Draskoviceva 13, HR-10 000 Zagreb, Croatia
E-mail: mjelavic@yahoo.com